

RECOMMENDATION ITU-R P.371-7

CHOICE OF INDICES FOR LONG-TERM IONOSPHERIC PREDICTIONS

(1963-1970-1974-1978-1982-1986-1990-1995)

The ITU Radiocommunication Assembly,

recommends

- 1 that the 12-month running mean sunspot number R_{12} be adopted as the index to be used for all ionospheric predictions for dates more than twelve months ahead of the date of the last observed value of R_{12} ;
- 2 that R_{12} , or the 12-month running mean value of Φ , the 2 800 MHz solar radio noise flux, (i.e. Φ_{12}) be adopted as the index to be used for predicting monthly median values of foF2 and M(3000)F2 for dates certainly up to six, and perhaps up to twelve months ahead of the date of the last observed values: substantially equivalent results should be obtainable by the use of either of these indices;
- 3 that Φ_{12} be adopted as the index to be used for predicting monthly median values of foE and foF1 for dates certainly up to six, and perhaps up to twelve months ahead of the date of the last observed value;
- 4 that caution be shown in the use of the recommended indices at high magnetic latitudes, where the resulting ionospheric predictions may not be sufficiently accurate.

ANNEX 1

1 Introduction

The concept of indices for long-term ionospheric predictions relies on the assumption that the important characteristics of the ionosphere, such as the critical frequencies of the various layers and the MUF factor M(3000)F2, depend in a systematic way on certain measurable quantities concerned with solar radiation. It should however be noted that the correlation between these indices and the actual ionospheric characteristics does not necessarily imply a causal relationship, but rather an indication of associated phenomena. Changes of solar activity in general contain three components:

- a fairly regular component with a period of about 11 years, which represents the well known cycle of solar activity;
- a component that has a quasi-period of about a year or a little less; and
- erratic fluctuations with periods of less than a month.

2 Sunspot numbers

For studies of the main component of the solar cycle the 12-month running mean sunspot number R_{12} is used because the resultant smoothing considerably reduces complicated rapidly-varying components but does not obscure the slowly varying component.

The definition of R_{12} is:

$$R_{12} = \frac{1}{12} \left[\sum_{n-5}^{n+5} R_k + \frac{1}{2} (R_{n+6} + R_{n-6}) \right] \quad (1)$$

in which R_k is the mean of the daily sunspot numbers for a single month k , and R_{12} is the smoothed index for the month represented by $k = n$.

The two main disadvantages in the use of R_{12} are:

- the most recent available value is necessarily centred on a month at least 6 months earlier than the present time;
- it cannot be used to predict the shorter-term variation in solar activity.

Nonetheless, R_{12} appears to be the most useful parameter for long-term studies and predictions concerning the F2 layer.

3 Index Φ

Consistent and reasonably long series of observations of the solar radio noise flux at about 10 cm wavelength have been made by Canadian, Japanese and other laboratories. The monthly mean, Φ , of the daily values at Ottawa, expressed in units of $10^{-22} \text{ Wm}^{-2} \text{ Hz}^{-1}$, should be regarded as the reference data for this index. Φ is more closely correlated with E-layer critical frequency than are noise flux values at other wavelengths. As the solar flux observations are only available from 1947, the sunspot numbers remain one of the longest series of observations of a natural phenomenon. Therefore, the continued collection and recording of sunspot observations is encouraged.

4 Other indices

An index I_{F2} has been developed, based on measurement of foF2 at noon at long-established ionospheric observatories. A study reported in 1964 showed that I_{F2} was more closely correlated with foF2 than any other index available at that time. An index IG , a variant of the index I_{F2} , was developed (now discontinued) for use with the ITU-R maps for foF2 contained in Recommendation ITU-R P.434. A further index, T , developed in Australia uses measured data from 30 stations and is based on the mean of the monthly median values of foF2 at each hour. Apart from the use of data from all hours, the derivation of the index is similar to that for I_{F2} .

5 Correlation between various indices

The following figure and equations show the dependence of Φ and Φ_{12} on R_{12} .

Figure 1 shows the mean relationship between R_{12} and Φ as determined from data for the years 1947-1966. This corresponds to the expression:

$$\Phi = R_{12} + 46 - 23 e^{-0.05 R_{12}} \quad (2)$$


The recommended relationship between R_{12} and Φ_{12} , also evaluated in Fig. 1 is:

$$\Phi_{12} = 63.7 + 0.728 R_{12} + 8.9 \times 10^{-4} R_{12}^2 \quad (3)$$

6 The prediction of indices

There is as yet no method whereby it is possible to predict accurately indices for the next sunspot cycle, or, more generally, for a cycle which has not yet begun. Indices that have been calculated by using harmonic analysis or by using empirical and statistical laws applied to observations over some earlier and even recent cycles, have not proved useful in predicting those for a new cycle. After a sunspot minimum has been observed, future development of the cycle can be extrapolated to a certain extent, although the deviations have been observed to be rather extreme.

FIGURE 1
 Relationship between R_{12} and Φ or Φ_{12}



D01

In the United States of America R_{12} is predicted using an improvement of the McNish-Lincoln objective method. First a mean cycle is computed from all past values of R_{12} starting from the sunspot minimum of each cycle through eleven years thereafter. For prediction of a value in the current cycle the first approximation is the value of the mean cycle at the stated time after minimum. This estimate is improved by adding a correction proportional to the departure of the last observed value for the current cycle from the mean cycle. With the current computer programs a new prediction for each month of the remainder of the cycle can be made as soon as a new observed value becomes available. The statistical uncertainty of the prediction is fairly small for the first few months after the last observed value but becomes large for predictions 12 months or more in advance. As soon as a minimum is identified, new correction factors can be computed by including the observed values for the preceding cycle, for application to the new cycle.

Predictions of R_{12} for one year ahead are also carried out by the Sunspot Index Data Centre (SIDC) in Brussels. Their predictions for solar cycle 22 are shown in Fig. 2 and can be compared with the observed smoothed values.

Predictions of Φ_{12} , based on the McNish-Lincoln method, are carried out by the Radiocommunication Bureau (BR).

Measured and predicted values of R and Φ and their 12-month running mean values (R_{12} and Φ_{12}) are published by the BR in the monthly Circular of Basic Indices for Ionospheric Propagation.

The SIDC also makes available, via electronic mail, the measured and predicted values of R , with access through the File Transfer Protocol anonymous procedure.

FIGURE 2
Predicted and observed sunspot numbers, R_{12}

