



International Telecommunication Union

# Clinical Study on MPEG-2 Compression Based on the Rec. ITU-R BT. 500-9

Isao Nakajima, M.D., Ph.D.

Associate Professor

Tokai University Institute of Medical Sciences

Invited Researcher

Communication Research Laboratory, Japan

Workshop on Standardization in E-health  
Geneva, 23-25 May 2003



ITU-T

# Agenda

## 1, Useful ITU Recommendation !

Recommendation ITU-R BT. 500-9

< suitable data speed of MPEG-2 compression >

## 2, Application

Ambulatory applications

Real time telemedicine

< data speed x traffic numbers = bandwidth >



ITU-T

# User report on ITU Recommendation Clinical Study on MPEG-2 Compression Based on the Rec. ITU-R BT. 500-9


## Purpose

This research is intended to identify the optimum transmission rate of clinical-use MPEG motion pictures by the **Double-Stimulus Continuous Quality Scale** method based on the Recommendation **ITU-R BT. 500-9**.




ITU-T

## Frame Structure of MPEG & H.261



time  $\Rightarrow$



MPEG-1

MPEG-2

I : Intra Picture

P: Predictively Picture

B: Bidirectionaly Picture


Motion JPEG

H.261( Video conference )



ITU-T

## Motion Compensation






ITU-T

## Quality degradation of MPEG images

The following are examples of types of degradation in image quality caused by MPEG compression.



**Block noise**

1. Block noise
2. Mosquito noise
3. Color smearing




**Color smearing**



ITU-T

# MPEG-1 power density tested with DFT 3.0 Mbps v.s. 0.5 Mbps

Power Density Distribution of Synthesis Signal from MPEG-1  
Fast Fourier Transform ( Discrete Fourier Transform )





ITU-T

## Display Protocol ( ITU-R BT.500-9 )

1. Gather 10 examiners who are not familiar with image compression.
2. Explain to the examiners the test contents, presentation sequence, and how to fill out the evaluation form.
3. Using an image that is not used in the test, demonstrate quality degradation, features of degradation, and degree of degradation.
4. Request that the examiners point out clinical problems associated with image degradation.





ITU-T


## Display Protocol ( ITU-R BT.500-9 )

5. Display the original image followed by a compressed image, twice each in this order, and have the examiners fill out the evaluation form immediately after the completion of each presentation.
6. The order of the displayed images is not changed until all examiners have completed the test.
7. The first five images are dummy pictures (examiners are not informed of this), and the evaluations of those images are not included in the test results.
8. The test time must not exceed 30 minutes per examiner.



ITU-T


## Evaluation System





ITU-T

## Double Stimulus Continuous Quality Scale Method Presentation Structure of Materials



A: Original  
B: MPEG-1  
G: Mid grey



ITU-T




Mid gray scale



Mid gray scale


ID, NO:  
SEX: AGE:  
D. O. BIRTH:  
98/03/04  
10:43:01  
SCU——1



COMMENT:

Original

ID, NO:  
SEX: AGE:  
D. O. BIRTH:  
98/03/04  
10:43:01  
SCU——1




COMMENT:

Compressed



ITU-T

## Double Stimulus Continuous Quality Scale Method Presentation Structure of Materials



A: Original  
B: MPEG-1  
G: Mid grey



ITU-T

## Quality-Rating Form of the DSCQS

Picture No.	10		11		12	
	A	B	A	B	A	B
Excellent 4	<input checked="" type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Good 3	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Fair 2	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Poor 1	<input type="radio"/>	<input checked="" type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Bad 0	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

N. Narita: "Subject-evaluation Method for Quality of Code Images"







IEEE Trans. Broadcast., BC-40, 1, pp. 7-13 ( Mar. 1994 )

Workshop on Standardization in E-health





ITU-T

## Digital Video Format



ITU-R BT.601  
Format MPEG-2



**SIF**: H.261, MPEG-1

**SIF**(Source Input Format)




ITU-T

## Test Results

### Comparison MPEG-2 and MPEG-1

4:2:2(720x480)      SIF(360x240)







ITU-T

## Comparison as with frozen picture



Original



6 Mbps




4 Mbps



3 Mbps



1.5 Mbps




1 Mbps



ITU-T

## Test Results Comparison 2H and 6H

PVD ( Preferred Viewing Distance, H:Screen Height )






**ITU-T**

## **Conclusions**

1. The Double-Stimulus Continuous Quality Scale method based on the Recommendation ITU-R BT. 500-9. is the better methods to find small distortion on MPEG compression.
2. PVD 2H shall be used in routine clinical use, and PVD 6H for educational purposes and the detection of lesions.
3. Video endoscopy and CCD image compressed with MPEG-2 shall be required at least 3.0 Mbps.
4. The SIF image compressed with MPEG-1 shall be required 1.5 Mbps.



# **Video Transmission**


## **What we are aiming to !**

- 1. Problem of 3G(IMT2000) !**  
**Frequency selective fading**  
**Blocking**
- 2. How we can track a satellite from a high speed vehicle.**
- 3. IP-based multimedia for ambulatory applications.**
- 4. We can reduce medical cost!!**




ITU-T

# What is DPC via telecommunications ?





# Target images



## Video Endoscopy( Air way )



## Light reflex



## Video Endoscopy( Stomach )





**Abd. Echo**



**Frequency selective fading due to multipath**

# Flooring Phenomenon

Error ratio can't improve with high speed vehicle  
under the environment of multipath











## IMT-2000(3G)

2Mbps data up-link from moving vehicle is impossible with IMT-2000 (3G).


Only 144Kbps expected, but no hardware !!



# Blocking ( Shadowing )




# Elevation > 80 deg. Duration [ hour ]





TCR 15:12:26.18  
PLAY  
REMAIN 52 min

Visibility study ( natural image )





Natural image ? Binary image ? Analysis



ITU-T

# Result of visibility study

diversity distance shall be 2.3m





ITU-T



# Super Ambulance

with diversity satellite antenna(distance 2.3m)








# How to track a communication satellite from high speed vehicle




## Quadrant detector




## Non-backlash






N-star(GEO) tracking with Quadrant Detector

# Real-time Telemedicine MPEG-2, -4, 6 (3+3)Mbps




ID, NO:  
SEX: AGE:  
D. O. BIRTH:  
98/03/04  
10:43:01  
SCV——1

NAME:



COMMENT:





ITU-T

# Reduce Medical Cost !

## Medical Cost

In case of Acute Myocardial Infarction in Japan,  
Thrombolytic agents are effective within 60 minutes after the  
attack. To make fine diagnosis of AMI, we have to send 12-lead  
ECG and Echography from moving vehicle at real time.

If 50% of patients will benefit => reduce medical cost  
**=> over U.S. \$ 2 Billion/10 years**

## Satellite System Cost

Total cost of the Quasi-Zenith satellite system  
( 3 space crafts, 4800 ground terminals, TTC & operation)

**=> less than U.S. \$ 2 Billion/10 years**



ITU-T

## Summary of trials

- 1, Video in an ambulance shall effective to support real-time telemedicine.
- 2, 3G ( IMT-2000 ) cannot afford because of its limitation of the data speed.
- 3, IP-based multimedia for ambulatory applications can save human life and reduce the medical cost.
- 4, We need challenge !!

# Thank you !

This study was performed through special coordination funds of the Ministry of Education, Culture, Sports, Science and Technology, the Japanese Government. Thanks to Prof. Leonid Androuchko ( ITU-D SG-2 Q14/2 Telemedicine ) and Mr. Yasuhiko Kawasumi ( Q10/2 Rural communications ) for their many useful suggestions.

