BE HE@LTHY BE MOBILE

A handbook on how to implement **mHypertension**





A handbook on how to implement mHypertension

ISBN 978-92-4-000144-2 (WHO)

ISBN 978-92-61-29851-7 (ITU)

© World Health Organization and International Telecommunication Union, 2020

Some rights reserved. This work is available under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 IGO licence (CC BY-NC-SA 3.0 IGO; https://creativecommons.org/licenses/by-nc-sa/3.0/igo).

Under the terms of this licence, you may copy, redistribute and adapt the work for non-commercial purposes, provided the work is appropriately cited, as indicated below. In any use of this work, there should be no suggestion that WHO or ITU endorses any specific organization, products or services. The unauthorized use of the WHO or ITU names or logos is not permitted. If you adapt the work, then you must license your work under the same or equivalent Creative Commons licence. If you create a translation of this work, you should add the following disclaimer along with the suggested citation: "This translation was not created by the World Health Organization (WHO) or the International Telecommunication Union (ITU). Neither WHO nor ITU are responsible for the content or accuracy of this translation. The original English edition shall be the binding and authentic edition".

Any mediation relating to disputes arising under the licence shall be conducted in accordance with the mediation rules of the World Intellectual Property Organization (http://www.wipo.int/amc/en/mediation/rules).

Suggested citation. A handbook on how to implement mHypertension. Geneva: World Health Organization and International Telecommunication Union, 2020, Licence: CC BY-NC-SA 3.0 IGO.

Cataloguing-in-Publication (CIP) data. CIP data are available at http://apps.who.int/iris.

Sales, rights and licensing. To purchase WHO publications, see http://apps.who.int/bookorders. ITU Publications can be obtained from ITU Bookshop http://www.itu.int/en/publications. To submit requests for commercial use and queries on rights and licensing, see http://www.who.int/about/licensing.

Third-party materials. If you wish to reuse material from this work that is attributed to a third party, such as tables, figures or images, it is your responsibility to determine whether permission is needed for that reuse and to obtain permission from the copyright holder. The risk of claims resulting from infringement of any third-partyowned component in the work rests solely with the user.

General disclaimers. The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of WHO or ITU concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted and dashed lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by WHO or ITU in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by WHO and ITU to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall WHO or ITU be liable for damages arising from its use.

Printed in Geneva, Switzerland

BE HE@LTHY BE MOBILE

mHypertension

A handbook on how to implement

Contents

Acknowledgements Executive summary Background Rationale for mHypertension What is an mHypertension programme?

Developing an mHypertension programme





Operations management Content development and adaptation





Technology specifications

Monitoring and evaluation

Conclusion

References

Annex 1: Examples of mHypertension programm individual and system level

Annex 2: List of behaviour change techniques in mHypertension programmey

Annex 3: The mHypertension Message Library

Annex 4: Examples of mHypertension programm

Annex 5: Examples of evaluation questions

06
07
08
00
10





Promotion and recruitment

	49
	50
nes at the	52
ncluded in the	56
	57
me algorithms	58
	61

Acknowledgements

WHO and the International Telecommunication Union (ITU) gratefully acknowledge the following contributors to this handbook

Handbook coordination

WHO Department for Management of Noncommunicable Diseases, Disability, Violence and Injury Prevention: Cherian Varghese, Taskeen Khan and Hongyi Xu

WHO/ITU Be He@lthy, Be Mobile team: Melissa Harper Shehadeh, Briana Lucido, Allison Goldstein, Sameer Pujari, Virginia Arnold, Vinayak Prasad, Per Hasvold, Surabhi Joshi, Natalia Wroblewska, Evan Pye, Hani Eskandar and Simona Pestina

Content development

Kirsty Bobrow (Global Brain Health Institute), Donald DiPette (University of South Carolina), Elizabeth Dunford (George Institute for Global Health), Georg Ehret (Geneva University Hospitals), Alireza Mahdavi Hezaveh (Ministry of Health and Medical Education, Islamic Republic of Iran), Andrew Moran (University of Columbia), Leila Pfaeffli Dale (University of British Columbia), Devashetty Praveen (George Institute for Global Health) and Ajay Vamadevan (Centre for Chronic Disease Control), Paul Whelton (Tulane University School of Public Health and Tropical Medicine)

Guidance

Be He@lthy, Be Mobile Steering Committee members: Nick Banatvala, Douglas Bettcher and Edward Kelley, WHO; Doreen Bogdan, Stephen Bereaux and Marco Obiso, ITU

Further contributions

Prebo Barango (WHO Regional Office for Africa), Marcelo D'Agostino (WHO Regional Office for the Americas), Heba Fouad (WHO Regional Office for the Eastern Mediterranean), Clayton Hamilton (WHO Regional Office for Europe), Mina Kashiwabara (WHO Regional Office for the Western Pacific), Jagdish Kaur (WHO Regional Office for South-East Asia), Kathleen Lannan (WHO Regional Office for the Western Pacific) Ahmed Mohamed Amin Mandil (WHO Regional Office for the Eastern Mediterranean), Mohamad Nour (WHO Regional Office for the Eastern Mediterranean) and Carrie Peterson (WHO Regional Office for Europe)

Administrative support

Florence Taylor (Be He@lthy, Be Mobile team)

Editing Teresa Lander

Layout and design Blossom | blossoming.it

Executive Summary

WHO and the International Telecommunication Union (ITU) have formed a partnership to use mobile technology to help combat noncommunicable diseases. As part of this Be He@lthy, Be Mobile initiative, WHO and ITU aim to assemble evidence-based and operational guidance to assist countries and governments to implement mHealth programmes for noncommunicable diseases. Mobile health, or mHealth, is defined as "medical and public health practice supported by mobile devices, such as mobile phones, patient monitoring devices, personal digital assistants, and other wireless devices" (1). The Be He@lthy, Be Mobile initiative generally uses basic technologies common in most mobile phones. The handbooks specific to the mHealth programme act as aids to policy-makers and implementers of national, or large-scale, mHealth programmes.

This handbook provides guidance for developing and implementing an mHealth programme to support people with hypertension in achieving better control of their condition



and improving their quality of life. It describes how an mHypertension programme can be used to supplement and augment existing national hypertension control programmes. and illustrates the steps required for successful implementation. The example text message library provided in the handbook uses evidencebased behaviour change techniques to support healthy behaviour changes and hence better hypertension control. This document and programme content are informed by a relevant evidence base and were developed following standardized processes. These mHealth interventions have shown promise for the behaviour change necessary to reduce the risk of developing or exacerbating hypertension, and WHO considers that the proper application of this handbook can be a critical factor in success in achieving behaviour change.

The handbook provides guidance on the following components of an mHypertension programme:

Background

Hypertension, or blood pressure equal to or greater than 140/90 mmHg (2) is the leading risk factor for premature mortality globally and is estimated to cause 9.4 million deaths every year, which equates to more than half the approximately 17 million deaths due to cardiovascular disease (3). Blood pressure level is directly associated with cardiovascular risk and overall morbidity and mortality risk; lowering the blood pressure reduces these risks. Evidence suggests that even modest reductions in blood pressure are important and linked with risk reduction. Lifestyle-related noncommunicable diseases are chronic diseases that take years or decades to manifest; preventing or even delaying the onset of these diseases could improve lives and result in substantial cost savings (4). Addressing behavioural risk factors such as an unhealthy diet, physical inactivity, tobacco use and harmful use of alcohol can help to prevent hypertension at a population level. Figure 1 below shows the main factors that contribute to the development of hypertension (2). WHO and the United States Centers for Disease Control and Prevention have launched the Global HEARTS initiative to strengthen prevention of cardiovascular disease globally (5). The HEARTS technical package includes recommendations to countries for preventing and managing hypertension, a major risk factor for cardiovascular disease. Early detection of hypertension is key to minimizing the chance of heart attack, heart failure, stroke and kidney failure. The HEARTS technical package provides information on the four main behavioural risk factors for cardiovascular disease (5):



Control of hypertension can be achieved by a healthy lifestyle and safe and effective medications, when needed. However, to do so demands multistakeholder collaboration, involving governments, civil society, academia, and the private sector. mHealth presents one tool which supports the implementation efforts for hypertension control aligned as part of the implementation of HEARTS.

FIGURE 1. MAIN FACTORS THAT CONTRIBUTE TO THE DEVELOPMENT OF HIGH BLOOD PRESSURE AND ITS COMPLICATIONS



RATIONALE FOR mHYPERTENSION

In 2016, ITU reported that, in countries with available data, 85% of the population aged 25-75 years owned a mobile phone (6). This worldwide mobile phone usage makes mHealth a widespread modality for providing public heath interventions. While mHealth research has mainly taken place in high-income countries, the explosion of mobile technologies in lower-income countries means that it is now being explored as an approach to support the development and strengthening of health systems.

Much of the human and social impact caused each year by morbidity and mortality related to noncommunicable diseases could be averted through population level interventions that are well understood, cost-effective and feasible (7). Results from recent systematic reviews have been quite promising in respect of the use of mHealth to manage chronic conditions, such as coronary heart disease (8, 9) and its risk factor, hypertension (10, 11). mHealth interventions have also shown promise for the behaviour change necessary to reduce the risk of developing or exacerbating hypertension, such as engaging in regular physical activity (12, 13). Deploying mHealth at the population level can be an effective way to achieve behaviour change, especially in lower and middle-income countries where mobile phone penetration is growing exponentially and risk of cardiovascular disease is high (2).

Source: (2).

What is an mHypertension programme?

There are many ways mobile technology can be used to prevent or control hypertension. Annex 1 provides further details on mHealth programme ideas at the individual or health system level. This handbook focuses on the individual level, to provide basic education, behaviour change strategies (see Annex 2) and medication adherence prompts to support people with hypertension in achieving better control.

A comprehensive mHypertension programme operating over a wide geographical area should aim to address the spectrum of disease of hypertension, the use of appropriate technologies and the needs and cultural norms of the population. The interventions should be embedded in the continuum of prevention and care, since mHealth can support services at all levels (see Figure 2).

FIGURE 2. THE mHYPERTENSION CONTINUUM OF **PREVENTION AND CARE**



Previous mHealth hypertension interventions have delivered education and supportive strategies to make healthy lifestyle changes and/ or adhere to guidance using text messaging, mobile applications and telemonitoring (10, 11, 14, 15). However, the present handbook describes a standard mHypertension programme for the individual using the simplest form of mHealth technology: text messaging. A recent review found mHealth interventions to be effective for prevention and management of cardiovascular disease, and suggested that

The goal of the mHypertension programme is to help people with hypertension to improve their blood pressure control through healthy behaviours and self-management.

To achieve this goal, the focus of the messages is will receive three modules (adherence, healthy eating and physical activity), with the option on education and supporting behaviour change, namely adherence to prescribed medication and of adding modules on tobacco cessation and/ monitoring of blood pressure, eating a diet low or harmful use of alcohol if the user is a smoker in salt and high in fruit and vegetables, starting or consumes an unhealthy amount of alcohol. or maintaining regular participation in physical Details on the content for each stream can activity, stopping smoking and limiting alcohol be found in Section 2 of the handbook. Each consumption. Text message content is based stream will also have slightly different evaluation on the recommendations from the HEARTS outcomes, which can be found in Section 5 of technical package (5) and the WHO global brief the handbook. on hypertension (2). Each mHypertension user

text message interventions were more effective than smartphone interventions. However, using multiple modalities should be considered for maximum outreach (16). The mHypertension message library and programme algorithms found in Annexes 3 and 4 have been developed for text messaging. Countries or groups can choose to adapt the programme algorithm, message content and technology they use based on the capacity, interests and needs of their respective populations.

Developing an mHypertension programme



FIVE AREAS OF THE mHYPERTENSION PROGRAMME



SECTION 1

OPERATIONS MANAGEMENT



1.1 mHEALTH NEEDS ASSESSMENT

A thorough mHealth needs assessment, including a readiness assessment, is a crucial step in developing an mHypertension programme. As with any large-scale intervention, it is essential to understand the context in which the intervention will be delivered. For the mHypertension programme to be effective, text messages need to be consistent with users' needs, their capabilities and impairments and their motivations. Messages can support self-management by incorporating relevant and effective behaviour change techniques, which provide strategies and encouragement to overcome barriers. Supporting a person diagnosed with hypertension to self-manage their condition may increase their understanding of hypertension and how it increases their risk of cardiovascular disease. It may also encourage the person to be an active participant in decisionmaking about their health and motivate them

The following information describes questions related to the goals and content of an mHypertension intervention that should be considered as part of a needs assessment.

Which behaviours contribute to hypertension? Which behaviours can control diagnosed hypertension?

Based on the HEARTS technical package (5), the four main behavioural risk factors for cardiovascular disease are the following:



Other considerations for the population include taking blood pressure medication as prescribed and receiving health-care service reminders, such as attending appointments with health workers and regular monitoring of blood pressure. Individuals should know their own blood pressure, especially if a close relative had or has hypertension, as this is another risk factor (2).

A good understanding of the most prevalent unhealthy behaviours in each country's specific context will help decision-making about the emphasis and frequency of the text messages in the mHypertension programme. Identification of these behaviours can be completed as part of the literature review during the needs assessment.

to engage in healthy behaviours (17). A needs assessment will help determine the relevant barriers and facilitators of behaviour change for people diagnosed with hypertension in each participating country's context.

A needs assessment also provides a vehicle for consolidating information for planning, identifying gaps in knowledge and helping with decision-making. Conducting a needs assessment will provide an understanding of the setting for the operations management teams. The needs assessment will involve visiting, observing and interviewing key informants and stakeholders and documenting existing resources. The results of the needs assessment will be useful in the development and implementation of a national mHypertension programme, and act as a baseline measure from which the programme can be monitored and evaluated.

What factors are going to help/hinder behaviour change?

Once the most prevalent and influential behaviours have been identified among the population, it will be important to understand key barriers and facilitators of change in these behaviours. This needs to be explored primarily from the mHypertension participants' perspectives, using a person-centred approach.

INTERNAL INFLUENCES

- □ Knowledge level
- □ Low risk perception
- Perceived stigma (e.g. overweight/ obese; ageism)
- □ Traditional care-seeking practices
- □ Myths and misconceptions
- Physical and psychological addiction
- □ Self-efficacy

EXTERNAL INFLUENCES

- □ Socioeconomic conditions
- □ Sociocultural norms
- Social stigma (e.g. overweight/ obesity; ageism)
- Access to quality and affordable health services

A needs assessment will be able to identify the extent to which and intensity with which these barriers influence behaviour change in the given context. A literature search should identify existing local studies and an exploratory qualitative study can identify such barriers and facilitators. Such a study would include conducting a set of interviews and/or focus group discussions with people with hypertension. It may also be useful to interview or conduct focus groups with family members and support people of those with hypertension. The sampling strategy for such a study would need to take account of age, gender, and whether they were diagnosed with hypertension.

What are the end users' digital literacy and habits?

As the "digital divide" has declined between different social groupings across the world, there may still be a considerable "digital use divide". This often reflects an underlying gap in digital literacy and digital engagement. Special consideration should be given to creating an inclusive programme which allows users with functional limitations, such as vision, hearing, or motor impairment, or those with low levels of digital literacy and engagement, to take part *(18)*.

Several tools are now available that can be used on a representative sample to appraise digital literacy and practices across a range of domains including operational, navigation, social, creative and mobile. For instance, the eHealth Literary Scale has been found to be a reliable and valid instrument for use in older populations *(19)*. Countries should consider validating such instruments in their own populations as part of the needs assessment.

The digital media needs assessment is critical to programme algorithm planning, as it will help to determine the number of messages sent, whether two-way messaging is used and whether voice messaging needs to be employed.

Table 1 lists additional considerations relating to the operation and implementation of an mHypertension programme that should be included in the needs assessment.

TABLE 1. CONSIDERATIONS FOR mHYPERTENSION NEEDS ASSESSMENT

ТНЕМЕ	CONSIDERATIONS
Current situation of hypertension	 Prevalence of hypertension Demographic data on nun hypertension, including a What are the priority popertension
State of national programmes for hypertension control	 mHypertension programs human resources, funding Approaches and algorith treatment and enhancing Existing or planned syner (e.g. cardiovascular disea Campaigns and health pr Involvement of nongover Relevant priorities of the Use of clinical manageme Gaps in medication supp equipment, etc., and any mHypertension
Target group	 Target populations Challenges and opportur Literacy and language co Mobile technology and o Functional ability issues i impairment) Knowledge levels, culture and behavioural trends responses
State of mobile communications	 Statistics on use of mobil consumers of text messa Penetration and use of sr Description of the mobile network providers and w including any related to h Whether unsolicited text and/or occurs Whether health services Cultural issues in the use The projected evolution particularly with respect smartphones Regulatory issues such as message transmission, re be sent each day Existing mHealth program messages

on

- umber and percentage of people diagnosed with age, gender and location (e.g. urban or rural) pulations and population segments?
- me objectives, constraints, institutional and g
- ms for identifying people in need of support, g adherence
- rgies with other chronic condition programmes ase, diabetes)
- romotion strategies for hypertension
- rnmental, community and patient organizations
- government and the health system
- ent guidelines
- ly, laboratory facilities, diagnostics, medical
- additional supplies needed to support
- nities within the population
- onsiderations
- other digital literacy considerations
- in the use of mobile phones (e.g. dexterity, visual
- al attitudes, perception of risk, current behaviour elating to hypertension and chronic conditions
- le phones and text messaging; costs to ging, data and calls
- martphones and mobile internet access
- e network environment (e.g. the number of /hether they provide "value-added services", health)
- messaging (spam) by companies is allowed
- use text messaging or smartphone apps of mobile phones
- of the mobile phone market in the near future, to increased penetration of smart- or semi-
- s spam, consent to receive programmes, cost of estrictions on the number of messages that can
- mmes that could incorporate mHypertension

Contextual, geographical, cultural and behavioural influences	 Individual and cultural attitudes to health care and self-management Determinants and risk factors of hypertension Cultural and social factors that may influence individuals from adopting a healthier lifestyle Community dynamics and actors that would support successful mHypertension outcomes Motivation and incentives for individuals to participate in an mHypertension programme People who can be considered champions, "trusted advisors" and how to leverage them Expected interactions of target populations with potential channels in health interventions: SMS (text message), web, app, phone call, interactive voice response, brochure, face-to-face coaching or consultation Convenience for the population Cost (who will pay: the population or the government?)
Stakeholders	 Relevant agencies, organizations, donors, companies, experts and their potential interest in supporting an mHypertension programme, including 1. national and subnational/regional levels of public health experts 2. funders of public health services 3. ministry of health and other relevant government departments and agencies 4. private and other health-care workers who work in hypertension 5. government agencies responsible for telecommunications (telecoms) and data protection 6. telecoms companies, mobile network providers and industry bodies or associations 7. any local mobile phone service providers or companies that provide mHealth services 8. patient associations; community advisory groups 9. health insurance companies; other private sector supporters 10. academic researchers (public health, hypertension, cardiovascular disease, behaviour change, social marketing, mHealth)
Promotion	 What channel will be used to promote the mHypertension programme How people will be registered on the programme What recruitment strategies will be used Incentives for participation Access to and availability of health-care services and resources
Further research	 Other areas in which further research is required to facilitate a successful pilot or "soft" launch and implementation; operational research.

1.2 PLANNING FOR INTEGRATION OF mHYPERTENSION WITHIN HEALTH-CARE SETTINGS

Integrating mHypertension within health-care settings offers an mHealth delivery mode to support hypertension control strategies for people diagnosed with hypertension. It requires buy-in from health-care providers who will likely be involved in promotion and recruitment of the mHypertension programme.

TABLE 2. FACILITATING IMPLEMENTATION AND UPTAKE OF THE mHYPERTENSION INNOVATION WITHIN TRADITIONAL HEALTH-CARE SETTINGS

Compatibility	Have we adapted mHypertension t approaches? And those of our com
Relative advantage	Will mHypertension help us to do t contact patients with results, supp faster progress towards the goal o
Level of complexity	Have we made mHypertension eas arrangements for training, support oversight?
Observability of results	How easily and quickly will mHype with users and countries? Have the different groups will understand an
Triability	Have we organized it so that every before having to decide about add among all stakeholders been allow

The mHypertension package will need to be customized so that it is aligned with the needs assessment, existing strategies, procedures, human resources and activities. For example, text messages that refer people to their health-care worker will need to provide correct information about where to go, what to do and whom to speak to. All those involved in the care or patient pathway should be trained and/or briefed prior to the pilot launch, and staff should be trained how to use and integrate the programme and refer and support their patients in the programme.

1.3 PROGRAMME LEADERSHIP AND PARTNERSHIPS

To facilitate planning, implementation and monitoring and evaluation of the mHypertension programme, a management team should be established, with clear responsibilities and accountability for the programme (see Figure 3):

 an international technical advisory group of experts to advise and support the in-country project team; The key is to determine how mHypertension can best be integrated into the existing or future operations and how it could improve outcomes for all involved. The mHypertension programme should supplement, not replace, self-management advice and instruction from health-care providers. To facilitate uptake, it may be useful to consider the five characteristics that have been demonstrated to influence the likelihood of adoption of a technological innovation, outlined below in Table 2 (20).

o fit well with our goals, objectives, operations, munities?

hings better, such as making it easier to ort, reminders, etc.? Will it help us to make f better hypertension control?

sy to understand and use? Have we made good , troubleshooting, project management and

rtension produce results that can be shared results been organized in a format that Id relate to?

vone can comfortably try out mHypertension opting it? Has enough time and consultation ved for the pilots?

- a national steering committee for overall direction and agreement;
- a national technical advisory group of in-country leaders to manage operations, content development and adaptation, promotion and recruitment, technology specifications and monitoring and evaluation.

FIGURE 3. PROPOSED STRUCTURE OF AN mHYPERTENSION MANAGEMENT TEAM

Promotion

Management

programme

of recruitment,

and dissemination

communications. marketing

WHO, ITU, and the Informal Expert Group

of public health and mHealth experts to assist in drafting the handbook and advising on implementation

International mHypertension **Steering Committee**

with representatives from the ministries of health and telecommunications and national and international representatives of WHO and ITU, to decide on overall direction and agreement

National technical advisory group

Government sectors (including health, telecommunications, business, media, treasury and planning) to set up the legal, technical and financial framework for a sustainable programme. This group will maintain links with a large group of potential partners such as the telecoms and software industry, local telecoms and mobile network providers, nongovernmental organizations, health professionals, academic and research organizations, health insurance groups, health service providers, civil society groups, opinion leaders, the media and others as appropriate

Operations

Management of overall programme operations, including needs assessment, work-plan, budget and legal aspects

Content

Development and adaptation of programme content

Monitoring and evaluation

Management of the development and implementation of monitoring and evaluation plans

Technology

Management of technical aspects of programme development and implementation

WHO, ITU, AND THE INFORMAL EXPERT GROUP

The international experts that assisted in drafting the country handbook also have experience in programme implementation and will be available to advise on technical aspects, legal issues, choice of platforms for scaling-up, and feasibility. Experts from international information technology organizations, health economists to governments and international agencies

and business development experts can also be invited to advise on models for programme sustainability.

INTERNATIONAL mHYPERTENSION STEERING COMMITTEE

A steering committee should be responsible for presenting the mHypertension programme to disseminate activities, share lessons and issues. The technical advisory group will assist engage the population and policy-makers. the national operational team in making decisions Representatives from all government sectors about the target population, type of programme, involved in the programme should be part of the programme objectives and design, promotion of steering committee. In addition, representatives the programme and the evaluation plan. from WHO and ITU headquarters should also be **PROJECT LEADERS FOR NATIONAL** members, contributing to decisions that maintain **OPERATIONS, CONTENT, PROMOTION,** the coherence with the Be He@lthy, Be Mobile **TECHNOLOGY AND MONITORING AND** initiative and sharing lessons between countries.

NATIONAL TECHNICAL ADVISORY GROUP

These groups include the people who will develop or adapt the programme for cultural relevance The technical advisory group should support, and technical accuracy, promote the programme, inform and advise the programme throughout operationalize and maintain it, integrate it into its inception, development, implementation and the health system and health promotion services evaluation. It should include the people necessary and evaluate it. They may include hypertension to make decisions related to funding and planning, and public health specialists, evaluators and those who will be involved in implementing, statistical experts, health promoters, behaviour promoting and evaluating the programme, change and communications experts (including those who can contribute to the long-term linguists) and consumer groups. These teams sustainability of the programme, and someone will report to, or be a subset of, the technical who understands or represents the mobile advisory group. network environment in the country. Regular meetings should be held for information-sharing **1.4 WORKPLAN DEVELOPMENT** and updates on progress. The technical advisory group will assign roles and responsibilities to This section provides a checklist or template various organizations in the different phases for developing a project workplan for use of development, adaptation, implementation, evaluation and continuing service provision. and adaptation by any country intending to implement an mHypertension programme (Table They should encourage discussion on overall programme ownership, funding and contracts or 3). Each section of the workplan is described in detail throughout the handbook. agreements on dealing with technical and other

TABLE 3. CONTENT CHECKLIST OF A WORKPLAN FOR AN **mHYPERTENSION PROGRAMME**

CONTENT CHECKLIST

Background and context

In this section, information from the needs assessment that applies to decisions on the implementation strategy is used.

- Present situation or context of hypertension treatment and care in the country
- Problem statement: a description of the problem of preventing and controlling hypertension
- National or government commitment to this project
- Process used in project identification or formulation, the information used and the stakeholders involved
- Relation to previous and current programmes or activities in hypertension care

EVALUATION

Operations management

In this section, planning decisions are made, a description of the programme to be implemented is created, and an operations management plan is developed that includes identifying who will be responsible for implementing the project and for ensuring provision of services.

- Overall project objective: for example, "to adapt the mHypertension programme for the population of X, particularly targeting Y people, and to implement it as a free national service"
- □ How the programme fits into national or regional strategies
- □ Strategy for operationalizing and promoting the programme
- How the initiative will evolve with progress in technology; become integrated into a comprehensive mHypertension and noncommunicable disease model
- Roles and responsibilities in the project: project team members, responsibilities for main activities; assign a project team leader
- Overall description of project management
- Management committee (if applicable, terms of reference)
- Accountability for project implementation

Public-private partnership

In this section, every partner has a unique role and all are motivated by the desire to improve health by using technology. Some important underlying principles to be considered include the following.

- A "win-win" philosophy, particularly for a long-term strategy, driven by the government, with the private sector and nongovernmental organizations involved in implementation
- □ A service free of charge to consumers to maximize public health impact (enrolment is very low when consumers have to pay)
- Consideration of the environmental effects of mobile networks
- Appropriate contractual arrangements with the best providers
- □ Assurance that service provision is sustainable in the long term

Content development and adaptation

In this section, a research driven message refinement process should be implemented.

- Review of existing programme content and rules of implementation
- Refinement of content through focus groups and consumer pre-testing, translation, changing the wording, removing and adding new messages, reframing messages, changing the rules in the system, and designing the registration, opt-in/opt-out and administration functions of the new programme
- Plan for updating messages

Promotion and recruitment

In this section, decisions should be made about marketing of and enrolment in the programme.

- Promotion and recruitment and retention plan (launch, short-term, mid-term and long-term) strategy)
- Recruitment methods (by SMS, web, missed calls, third parties)
- Promotion strategy (media, health workforce training, civil society outreach, etc.) adapted for various client groups (by demographics, e.g. urban, rural, age, gender, income)

Technology

In this section, decisions are made about the considerations necessary for the infrastructure and rules of the programme.

- □ The type of mHealth technology and channels to be used (SMS, MMS, voice, apps, etc.)
- Availability of technology options within the public sector and/or private sector
- Process for procurement and adaptation of technology
- Dashboard development and access
- Procurement of a short code
- Negotiation with telecoms regulators, aggregators and operators for pricing
- Data security
- Technology pre-testing and scale-up plans

Monitoring and evaluation

In this section, decisions should be made about what will be measured by the programme, by whom and with what frequency.

- Monitoring and evaluation plan based on the mHealth monitoring and evaluation framework
- □ Short-term, mid-term and long-term plans

Estimated time frames

Planning: 3-4 months

- Needs assessment: assuming the information is readily accessible and local researchers are available to inform the technical advisory group
- Formation and planning of the technical advisory group: engaging the right partners, agreeing on the implementation plan, the promotional plan and the evaluation plan; working groups formed; project leader assigned
- Content development team formed
- Selection of programme and text message content or preparation of new text messages
- Agreement on programme and technical and transmission specifications

Content adaptation: 4-6 weeks

- Expert group consultation to review and culturally adapt content
- Consumer testing of messages before use in focus groups, online surveys, telephone interviews, etc.
- Modification of messages based on consumer feedback
- □ Translation (and back-translation to check for translation accuracy)

Technology adaptation: 4–6 weeks

Development of promotion and recruitment strategy and material: 4-6 weeks

Programme implementation: within six months of beginning the planning

Monitoring and evaluation

- Monitoring: should be done in real time, from the start of programme planning until the end of programme implementation, using agreed indicators
- Evaluation: beginning, at least, with a six-month follow-up of the initial cohort

Reports and dissemination plans for evaluation, refinement, improvement and service provision

SECTION 2

CONTENT DEVELOPMENT AND ADAPTATION



(a) be informed by the existing evidence on effective mHealth interventions;

(b) be informed by the findings of a needs mHypertension (see Figure 4). assessment; and

FIGURE 4. DESIGNING A TEXT MESSAGING PROGRAMME



Table 4 lists key decisions for consideration by the content development team. Each decision contains recommendations based on previous mHealth and hypertension research and is further described in detail below with supporting evidence. This mHypertension handbook provides recommendations for programme content and delivery; however, it is up to each country to determine how it wishes to develop their programme on the basis of its needs assessment. The full message library and programme algorithm can be found in Annexes 3 and 4.

TABLE 4. DECISIONS AND SUGGESTED ACTIONS FOR mHYPERTENSION PROGRAMME CONTENT DEVELOPMENT

DECISION	ACTIONS
Target audience	Target people with hype
Goals of the	Help people with hypert
programme	healthy behaviour and se
Select behaviour	Target: information, re
change techniques	techniques (see full list o

development of communication interventions for health behaviour change.

Recently, a consensus has emerged on these steps (21) which have been adapted for

- Choose the communication objectives (e.g. beliefs, attitudes, knowledge)
- Design the framework or algorithm for the programme (timing and frequency of

ertension, as informed by a health provider.

ension to improve their blood pressure through elf-management.

eminders, instruction and self-management of behaviour change techniques in Annex 2).

DECISION	ACTIONS
Design the framework of the programme (duration, message frequency, order, repetition, time, directionality, framing)	 The programme should last at least six months (see Section 2.3). Messages should start at a high dose for the first few weeks and then gradually decrease in frequency over time. Start with general informative messages, moving to self-management with specific behaviour change techniques, and then to maintenance of behaviour change in the later phase of the programme. We recommend grouping messages into modules, with each module targeting a specific behaviour. Allow users to choose when they receive messages, if possible. Offer two-way messaging. Messages should be clear and direct, offering practical and relevant. advice for older adults in simple language. Messages should emphasize the benefits of action (positive framing) over the consequences of inaction (negative framing).
Message components	Messages target the lifestyle advice from the HEARTS technical package. To encourage behaviour change, messages incorporate relevant behaviour change techniques.
Pre-testing, piloting, refining messages	Once a first draft of messages and the programme algorithm has been developed, pre-test the messages with the end user. Refine messages after pre-testing, then conduct a pilot programme. Refine messages and algorithm after the pilot programme prior to national scale-up of programme.

2.1 TARGET AUDIENCE AND **GOALS OF THE PROGRAMME**

2.2 SELECT BEHAVIOUR CHANGE TECHNIQUES

Intervening at an early stage is essential because the early detection of hypertension can significantly minimize the risk of developing cardiovascular disease. The mHypertension programme will target those who have been diagnosed with hypertension by a health provider to achieve better control of their hypertension through healthy lifestyle change and selfmanagement strategies. The programme may additionally target the general population and health workers.

As part of the needs assessment, the steering committee should have agreed on a set of target behaviours and specific determinants (barriers and facilitators) to behaviour change, through relevant literature reviews and consulting with experts. The next task is to agree on the behaviour change techniques required to modify these behaviours and make healthy lifestyle changes. As highlighted in Figure 5, a behaviour change technique that is likely to modify a particular behaviour has to influence its determinant.

FIGURE 5. BEHAVIOUR CHANGE PROCESSES WITH EXAMPLE OF **EXERCISE FOR HYPERTENSION CONTROL**



Source: (23). BP: blood pressure.

For example (see Figure 5), if the desired outcome is to control blood pressure, one behaviour that helps is to engage in regular aerobic exercise, which has the benefit of lowering blood pressure below pre-exercise levels for 2-3 hours postexercise (23). To encourage this behaviour, the self-efficacy or self-confidence to do regular aerobic exercise may be required. A behaviour change technique that can increase self-efficacy is to provide graded tasks, where easy-toperform tasks are prescribed first, followed by increasingly difficult, but achievable, incremental tasks. An example message targeting graded tasks that is ultimately aimed at improved mobility is shown below.

Messages in the message library (Annex 3) have Try to walk or do other exercise for 30 been paired with a corresponding behaviour minutes each day to stay healthy. If change technique to ensure the mHypertension you are new to exercise, start at a slow programme has a theoretical framework. pace and try to add a minute each day. Internet- and mobile-based interventions with a theoretical underpinning have been shown to be Annex 2 provides a list and description of more effective than those not based on theory behaviour change techniques that may be (26, 27). The combination of theory and the relevant based on the taxonomy of Michie et HEARTS technical package recommendations al. (24). The list was compiled from reviews of creates a usable framework to underpin an techniques used in mHealth intervention to mHypertension programme.

2.3 DESIGN THE FRAMEWORK OF THE PROGRAMME

DURATION OF THE PROGRAMME

Recent reviews of mHealth interventions have found no clear conclusions on the optimal length of programmes, because of the low number of randomized controlled trials and heterogeneity of mHealth interventions (8, 28). Research shows it can take six months of engaging in a new behaviour to move into the maintenance stage of behaviour change (29, 30). Two mHealth trials delivering text message-based interventions to adults (mean age ~60 years old, N=294) with coronary heart disease found most participants thought 24-weeks was the right length for a programme (31, 32).

SUGGESTION 1

The programme should last at least six months.

TIMING AND FREQUENCY OF MESSAGES

As with the duration of mHealth programmes, few systematic reviews drew strong conclusions on effective message algorithms (28). Despite this, a few reviews have made recommendations on the number and frequency of messages, although all agree that the area needs further investigation. Evans et al. reported that a higher dose of messages resulted in greater behaviour change but gave no indication on the optimal number of messages per week (33). One review found message frequency that is customizable or declines in frequency over time were more effective than fixed frequency (a low or high number of fixed messages) (34). Another systematic review found that at least one message every three days appeared effective in changing behaviours (8). Park et al. conducted a review on mHealth interventions to prevent and manage cardiovascular disease, finding that studies with the highest frequency of messages

improve clinical outcomes and behaviour change in cardiovascular disease (8, 25). The most commonly used behaviour change techniques for secondary prevention of cardiovascular disease included the following:

- information about health consequences
- goal-setting (behaviour)
- self-monitoring of behaviour
- social support (practical)
- prompts/cues.

(i.e. at least once daily) had the strongest effect on adherence to medication and behaviour outcomes (16).

SUGGESTION 2

Messages should start at a high rate for the first few weeks, e.g. once per day, and then gradually decrease in frequency over time, with at least one message every three days for the duration of the programme.

ORDER OF MESSAGES

There was little evidence in the literature specifying the order of messages. Maar et al. recommended placing positive content before negative content, which may be more relevant for individual message framing (35). Other behaviour change and exercise prescription programmes began with more introductory/general messages first which then become more specific (31, 32). For instance, the messages could be more informational for the first few weeks, followed by specific instruction and self-management techniques. Specific behaviour change goals should start slow and offer gradual progression as the programme continues. The final half of the programme can incorporate messages that focus on maintenance of these newly adopted behaviours and reinforce the long-term benefits.

SUGGESTION 3

Start with general informative messages, moving to self-management with specific behaviour change techniques, and then to maintenance in the later phase of the programme.

REPETITION OF MESSAGES

Multiple behaviour change is under-studied; however, a growing body of evidence points to the effectiveness of changing multiple rather than single behaviours (36, 37). There have been few studies examining the effectiveness of changing multiple behaviours sequentially as opposed to simultaneously, but one review suggests that both sequential and simultaneous multiple health behaviour change is effective (36).

SUGGESTION 4

We recommend grouping messages into modules, with each module targeting a specific behaviour. Please see Table 5 below and the message library (Annex 3) for further clarification. To avoid repetition and participant boredom all messages should be unique. Part of the content development phase is to determine if users would rather receive modules simultaneously or sequentially.

TIME OF DAY MESSAGES ARE DELIVERED

One systematic review suggested that interventions were most successful when the time of day messages are delivered was customizable (34). Allowing user choices for delivery was also important to the participants in mHealth studies for self-management of cardiovascular disease; however, most participants chose to receive messages in the time slot 9 a.m.-12 noon (31, 32).

SUGGESTION 5

Allow users to choose when to receive messages if possible. If tailored programmes are too costly, deliver messages at a time of day when participants are most receptive (e.g. midday: not too late or too early).

DIRECTIONALITY OF MESSAGES

Two reviews have found two-way messaging was more effective than one-way (16, 34). However, many one-way mHealth interventions have been effective in changing behaviours (38, 39).

SUGGESTION 6

Encourage two-way messaging if feasible.

MESSAGE FRAMING

Many studies on message framing have shown that positive framing is more effective and acceptable. Positive messages are benefit-oriented rather than consequence-oriented (40). In one study, positively framed messages resulted in greater behaviour change, as older adults preferred and remembered positive messages better compared with negatively framed messages (35, 41). It is important to note that message framing preferences can vary by population and should be tested in the pre-testing/piloting phase to determine if the population is more motivated by the potential benefits of behaviours ("regular exercise can control your blood pressure") or by reducing the consequences of inaction ("if you continue to smoke you increase your risk of heart attack or stroke").

In message framing:

Do use (35, 40):

- \checkmark positive and encouraging tone
- reminders
- \checkmark one idea per message
- correct spelling and grammar
- \checkmark direct, simple and concise language

Advice should be clear, practical, and realistic for the intended user (35). Specific content should be targeted to what people with or at risk of hypertension find relevant (42) and what behaviours they need to change, as tailored and targeted messages are more effective at changing behaviour (16, 34). To ensure that the language is at an appropriate level, literary testing of messages in the language concerned is recommended (e.g. Google web corpus or Lexile Analyzer®) (42). Muench et al. (40) found that individuals with less education were more likely to prefer the inclusion of an emoticon than those with more education. There were no preferences in terms of punctuation, such as the number of exclamation points or capital letters compared with messages with no visible emphasis.

SUGGESTION 7

Messages should emphasize the benefits of action (positive framing) over the consequences of inaction (negative framing). Messages should also be clear and direct, offering practical and relevant advice for older adults in simple language. It is suggested that message content and tone should be pre-tested with end users as part of the pre-testing and piloting phase.

2.4 MESSAGE CONTENT

This section describes some key topics that the mHypertension message library could address. The topics are based on recommendations from the HEARTS technical package (5) and the WHO global brief on hypertension (2). As mentioned, it is recommended tailoring messages to people diagnosed with hypertension. There are five streams recommended for the programme; Table 5 outlines the content for each stream.

Do not use:

- × authoritarian and stern tones
- × immediate commands
- × over-simplification
- very informal "textese" ("ur", "l8r", "b4")

TABLE 5. CONTENT OF mHYPERTENSION PROGRAMME

MODULE	DESCRIPTION	EXAMPLE MESSAGE	MODULE DESCRIPTION
Adherence	 Medication Take your blood pressure medications exactly as prescribed. Often medication is needed to control blood pressure. You may need to take more than one type of medicine. A fixed-dose combination medication, where two medicines are in a single pill, can ease logistics and reduces pill burden. Even if you do not feel unwell, it is extremely important to keep taking your blood pressure medicines as prescribed. Know the generic name and side-effects of your drugs. Consult your doctor if you have any side-effects from the medication. Take your medication as prescribed, every day, regularly. Don't stop or change your medication without consulting your doctor. Talk to your doctor before taking a new drug. Adherence to treatment is critical for blood pressure control. Practising a healthy lifestyle should be combined with taking medicines. Hypertension is manageable and controllable, but not curable. Know the red flag signs for hypertensive emergency: severe headache and loss of consciousness, chest pain, nausea and vomiting, dizziness, visual disturbance, racing heartbeat. Do not miss appointments with your health worker. If you feel unwell, do not delay – call or visit your health worker. 	It is important to take your medication as prescribed. This means taking it at the right times and not missing any doses. Keep on top of regular blood pressure screening. Your blood pressure should be less than 140/90. If it is high, see your doctor.	Healthy eating • Eat a heart-healthy did risk of overweight and high blood pressure, did Eat less than 1 level teat eating too much a pressure; eating too much a pressure; • eating too much a pressure; Imit falvourings th soy sauce, fish sau g. Imit falvourings th soy sauce, fish sau many canned fooc amount of them th * Eat a least 5 portions Imit salty rodes su pickles, salty snach many canned fooc amount of them th * Eat at least 5 portions I portion is equal to or banana, or 3 tat I starchy vegetables (these include pot other starchy root) • Eat fewer fatty or oily eat less fatty mean baked goods; I when cooking, u olive, sunflower, s animal fat or palm boil, steam or bake i remove the fatty i before cooking; i eat white meat (for meat to once or ty eat nuts, lequemes, who
	 Monitoring/screening Do not miss appointments with your health worker. If you feel unwell, do not delay - call or visit your health worker. Have your blood pressure checked at every visit. People with raised blood pressure may monitor their own blood pressure if an appropriate device is available and affordable. Have your blood sugar checked once a year. For people with diabetes, refer them to the mDiabetes programme. Ask for your family's help to live a heart-healthy life. Individuals can choose to live a healthy life through making healthy diet choices, increasing physical activity levels, avoiding tobacco use and avoiding harmful use of alcohol. Keep a healthy weight. If you are overweight, losing weight 		 Eat nuts, legumes, who Eat fish or other food seeds) at least twice a Avoid added sugar: choose fresh fruit a of sugary snacks chocolate; drink fewer soft d are high in sugar (flavoured milks and reduce sugar in yo) When buying food i sodium (salt), sugar, familiar (salt), sugar, familia

- healthy diet increases your and of heart attack, stroke, cancer and other diseases.
- g or less) of salt per day: major cause of high blood
- ting or cooking but, better when eating or cooking;
- n a lot of salt (for example, on, or stock cubes);
- fish or meat, salty cheese,
- ad contain salt, so limit the
- bles and fruits per day:
- nple, 1 orange, apple, mango of cooked vegetables; ount as one of your portions
- veet potatoes, cassava and es).
- ssed meat, fried food and
- ier vegetable oils such as soya or corn oil, instead of ut oil;
- nan fry;
- eat, including chicken skin,
- chicken) and fish; limit red ek at most.
- and foods rich in potassium. nega-3 fatty acids (e.g. flax
- egetables as snacks instead cookies, cakes, candy and
- soda and other drinks that juices, cordials and syrups, drinks); drink water instead; d coffee.
- ore, check food labels for prie content.

EXAMPLE MESSAGE

Prepare food using only a little bit of salt to help you and your family maintain normal blood pressure. Keep the salt shaker off the table!

MODULE	DESCRIPTION	EXAMPLE MESSAGE
Physical activity	 Physical activity protects your health and can help you feel better and happier. Insufficient physical activity increases your risk of having a heart attack, stroke, cancer and other diseases. Increase to 150 minutes of moderate physical activity per week, equivalent to brisk walking, spread throughout the week. Moderate physical activity should make you breathe slightly faster and make your heart beat slightly faster. Start slowly. You can engage in physical activity for as little as 10 minutes at a time, adding these up to reach 30 minutes a day or 150 minutes spread over the week. People can do moderate physical activity at any age - do as much as you can manage. 	Walk to work, walk to the shops, take the stairs, play with the kids or take the dog for a walk. It all adds up - build physical activity into your day.
Tobacco cessation (add-on)	 Using tobacco increases your risk of having a heart attack, stroke, cancer and other diseases. Do not start smoking. If you smoke, remember that quitting tobacco is the most important thing you can do to protect your health. Ask your health worker to help you quit smoking. Do not allow smoking in your home: second-hand smoke damages the health of your family and other people around you. Smokers could also be referred to an mTobaccoCessation programme, if implemented, for extra support. 	Do not be fooled! There is no such thing as a safe cigarette. Quitting is the only way to protect yourself from the health risks of smoking.
Harmful use of alcohol (add-on)	 Drinking alcohol can increase your risk of having a heart attack, stroke, cancer and other diseases. Overall, the best way to avoid the health risks of alcohol is to abstain. If you do drink alcohol, keep in mind that less is better. Avoid having more than two drinks on any one day and do not drink any alcohol on at least two days per week. Do not drink alcohol for "health" reasons. Do not use alcohol when you are: driving; operating machinery; pregnant or breast feeding; taking medications that interact with alcohol; living with a medical condition that is made worse by alcohol; having difficulty in controlling your drinking. 	Think of supportive people you trust, like friends, family, or your partner. Ask them to help you cut back on alcohol.

SPECIAL CONSIDERATIONS FOR SALT/ SODIUM

The HEARTS technical package, specifically the H module, has dietary advice that includes reducing salt, increasing fruit and vegetable consumption and limiting unhealthy fats and sugar. For an mHypertension programme, special consideration (or emphasis) should be given to decreasing salt/sodium intake, as this is a commonly-cited prevention strategy to reduce the burden of hypertension (7). Data from around the world suggest that the average sodium consumption is well above the minimal physiological needs, and in many countries, it is above the value recommended by the 2002 Joint WHO/Food and Agriculture Organization of the United Nations Expert Consultation of 2 g sodium/ day (equivalent to 5 g salt/day) (43). The high prevalence of hypertension in adult populations globally, and the clear benefit of reduced sodium in individuals with high blood pressure, means that reducing sodium intake is likely to be broadly beneficial to populations around the world. For example, modelling the impact of decreasing salt intake from the estimated global level of 9-12 g per day (g/day) to the recommended WHO maximum target of 5 g/day has demonstrated the potential to reduce the global stroke rate by 23% and the global cardiovascular disease rate by 17%. This would prevent about 2.5 million deaths every year (44, 45).

2.5 PRE-TESTING, PILOTING AND REFINING THE MESSAGES

Once a good plan for the mHypertension programme has been developed, it should go through iterative stages of pre-testing, pilot and

Once mHypertension is pre-tested and piloted, refinement. it should go through a refinement process addressing all key issues highlighted during PRE-TESTING THE TEXT MESSAGES the pilot, before its final launch. Messages may need to be reviewed and updated several The draft text messages should be pre-tested by times. The programme plan should identify a conducting interviews or focus group discussions person responsible for maintaining and updating with potential users. It is important that such the message library. Having some degree of interactions should include representatives from flexibility and keeping the programme fresh various subgroups within the target audience. may help to keep the audience engaged. Once During these interactions, the facilitator/ the programme is launched, someone on the interviewer should explain the purpose and team should also be receiving the messages in content of mHypertension to the participants real-time and monitoring them closely to ensure and provide them with the message library, that the correct messages are being sent asking them to rate the messages on content, on schedule.

clarity and persuasiveness. Testing should assess message comprehension, language, message framing and tone. It is important to consider the literacy levels of the target population, including health literacy. Messages that are unclear or inappropriate for the target group should be rewritten, either by or with the participants.

PILOT (ALL VERSIONS OF THE PROGRAMME IF POSSIBLE)

Once the messages are pre-tested and refined, the mHypertension pilot programme should be launched. Such a pilot could last for approximately 2-4 weeks and involve 25-50 users. This pilot should be evaluated using two sets of data: users' self-reported experience and their realtime engagement with the programme. Users' experience can be collected either through faceto-face/phone-based interaction or via webbased/internet-based surveys. The key questions to ask include the user's experience of taking part in the mHypertension programme; clarity, quantity, timing and frequency of messages; what was good about the programme and what was not; completion/non-completion of the programme; and any effect on their attitude or target behaviour(s). Users' real-time engagement can be evaluated using records from mobile phone companies (if available) of messages delivered and opened. Depending on the complexity of the programme, these data can provide information on the extent of clients' engagement. Measures used in the monitoring and evaluation phase of the mHypertension programme can also be tested during the pilot stage.

REFINE AND OPTIMIZE THE PROGRAMME

SECTION 3

PROMOTION AND RECRUITMENT



A nationwide or population-specific strategy to promote outreach and recruitment to the programme should be considered early in the planning stages, taking account of the following considerations.

- Target audience and access to media/promotional strategies.
- model is likely to direct the promotional campaign.
- Existing health programme promotion strategies and synergies: which organizations/notable personalities are currently involved in mass media campaigns for hypertension prevention and/ or self-management services and can those campaigns be linked or leveraged? Which mHealth programmes have previously been implemented in the area? Can lessons be learned about which promotional techniques are effective?
- How people will register with or sign up to the programme: directly by text message, online, by telephone, in person or through a third party (e.g. health-care provider).
- The amount of data to be collected at baseline to allow tailoring of the materials and ensure follow-up.
- The local mobile network environment: e.g. whether sending unsolicited text messages is allowed (this violates network codes of conduct in some countries). This may also be an important consideration in the potential effectiveness of or engagement in the programme, i.e. whether a population that has been desensitized to receiving unsolicited health-related text messages will be likely to read mHypertension text messages.
- Use of incentives to encourage participation.
- Effectiveness of mobile messaging (MMS) for outreach and promotion, leveraging the marketing and promotional campaigns of the stakeholder agencies, including technical partners, such as telecoms companies. This can enable campaigns where consumers can self-issue health messaging around this topic, i.e. a health promotion and/or or social marketing programme with posters and radio advertisements, through community settings, places where people meet or through social media, where mobile numbers are available for texting and automated health information can be sent to anyone who requests it.
- Timing of the campaign: consider a soft launch prior to starting the promotional campaign to ensure that all processes are working well before the official start date for registration.
- Cost of radio and TV advertising. This advertising can be an important method of promoting the programme but, in many countries, it can be very expensive. Initial underestimation of promotional costs is common and can be difficult to remedy later.

Who the public considers to be the "owner" of the programme and the data, as the operational

SECTION 4

TECHNOLOGY SPECIFICATIONS

The following technical aspects of an mHealth programme must be considered by the technical advisory group from the start, in collaboration with local partners:

- type of mHealth technology and channels to be used (SMS, MMS, voiceover internet, apps (e.g. messaging apps), etc.);
- platforms that allow video could be particularly useful for demonstrating content in action, such as exercise content or skill-building, video messages demonstrating correct strength training techniques or how to cook;
- availability of technology options within the public and/or private sector;
- process for procurement and adaptation of technology;
- dashboard development and access;
- procurement of a short code (the 5- or 6-digit unique number used to send and receive SMS to and from mobile phones, "send a message to 123123 to enrol in the programme");
- negotiation with telecoms regulators, aggregators and operators for pricing;
- data security;
- plans for technology pre-testing and scale-up.

It is important to note that mobile communications network environments differ from country to country. The specificities should be considered in the planning stage by including in the technical advisory group technical experts (such as representatives of telecoms companies, operators, telecoms regulatory authorities, government departments responsible for information, communication and technology) or individuals knowledgeable about the communications network in the country. Network operators, telecoms companies or industry organizations can provide help in setting up the programme and advising on its sustainability.

Some providers may consider supporting such a programme as good publicity or a useful addition to the services they offer. In the absence of such support, the programme can be delivered through a contractual arrangement with an "aggregator" or "gateway" company that has established relations with all telecoms companies and networks (see Figure 6). This can be a cost-effective way to deliver messages to many participants, regardless of their mobile carrier or location, without establishing these interfaces individually. Although the aggregator adds a further cost, this cost decreases as the scale of the programme increases; using an aggregator can therefore be more cost-effective than attempting these activities "in-house", unless the necessary capacity and infrastructure already exist.

FIGURE 6. TWO-WAY SMS SYSTEM WITH AGGREGATOR AND SHORT CODE



When thinking about the technical aspects of an mHealth programme, the technical advisory group may also consider the following questions.

- Partnerships (see Table 6): what sort of arrangement with telecoms companies and/ or the aggregator will best suit long-term implementation of the programme?
- Communicating messages (see Table 7): should voice messages, video messages, interactive messaging or interactive voice response systems be used? What are the capacity, cost-effectiveness and reach of the available technologies in the country?
- Free access: how can we ensure that the programme is free and available to all consumers regardless of their carrier, network or location?
- Market research: which telecoms system is most appropriate in the country, based on reach (subscribers), coverage, costs, security and sustainability? Who owns the data? What are the privacy regulations and how will data be protected and kept secure? Is there good interoperability with health systems? What are the considerations for data protection and how should a central database best be maintained?

- Sustainability: what are the operating costs of the programme, such as the cost per message, and how will these affect the scale of the programme?
- Contracts: in establishing contractual arrangements with partners, what are the considerations regarding intellectual property, security and privacy of mobile phone numbers, testing, expectations of involvement in monitoring and evaluation and service agreements? Who will manage the contractual arrangements, and what support will be given for maintenance and any other problems?

The technical advisory group should also consider logistics and the functional plan and finalize the functional specifications in collaboration with technical partners who will build the appropriate systems and interfaces and test internal and user acceptance. Table 6 provides an overview of the roles of the various stakeholders to be considered when planning the technological aspects of an mHealth programme.

TABLE 6. ROLES OF STAKEHOLDERS IN SETTING UP mHEALTH TECHNOLOGY

STAKEHOLDER	ROLE
Ministry of Health	 Official owner and custodian of the programme, part of the governance body Assess and identify needs, develop and validate content Contract service providers or build in-house infrastructure/platform Sign cooperation agreements with all operators and/or service provider Fund or partially fund the programme Can host the mHealth platform/database and own the short code Manage the promotion and marketing campaigns
Telecommunications ministry eGovernment (e-Gov) entity (if applicable)	 Policy-making to enable m-services in terms of regulations and policies (Partially) fund the programme, participate in the governing body Provide technical expertise to the ministry of health Possibly host the platform Facilitate dialogue between ministry of health and stakeholders in information and communications technology (ICT) Support the negotiation of preferential prices for m-services
Telecommunications regulatory authority	 Verify eligibility for short-code acquisition Allocate short code Facilitate dialogue between ministry of health and ICT stakeholders Fund or partially fund the programme
mHealth service providers (if ministry of health or eGov does not have a platform)	 Provide SMS management application/platform Manage the platform and run SMS campaigns Provide 24/7 technical support Deal with telecoms operators, potentially manage the short code
Telecoms operators	 Deliver SMS to end users Set the cost of SMS and agree on special tariffs with ministry of health, if possible Facilitate interfacing with service providers and/or local aggregator Support the promotion of the service
Local aggregator	 Provide interface with all operators and manage relationship and invoicing process Provide reporting on services delivered/failed (Potentially) own and manage the short code
Data privacy commission	 Set the rules for data protection Enforce the application of data protection regulations Authorize mHealth services, providing they respect data privacy Authorize data storage outside the country, if necessary
WHO and ITU	Provide technical expertise and share knowledge from other countriesHelp to convene stakeholders

TABLE 7. EXAMPLES OF TECHNOLOGY OPTIONS FOR HEALTH PROMOTION

CHANNEL	STRENGTHS
Interactive voice response (IVR)	 Voice- and phone-enal access Fast time to market Supports natural lange Ease of integration
SMS	 Simple, easy and conv Cost-effective Private communicatio Fast communication
Unstructured supplementary service data (USSD)	 Simple and logical Real-time, fast and res Inexpensive Interactive navigation
MMS	 Direct and personal Messages can be sinforwarded Interactivity through r
Data applications	 Self-contained experies Graphics and user-get content Automatic updates and readable offline Leverages device-national capabilities (camera, or positioning system) Strong paid model
Mobile web	 More economical than apps Mobile phones and smartphones support. Mobility for content a services Videos and photos

	WEAKNESSES
abled Juage	 Limited capability and development tools Inability to pause, resume, fast-forward or rewind
venient on	Some security vulnerabilitiesFake SMS (spoofing)
sponsive	 Session-based timeouts Codes more difficult to remember than common short codes
stored and multimedia	 Not compatible with basic phones More expensive than SMS Content adaptation limited by variations in screen size and resolution Read and response rates lower than SMS
ience enerated nd content tive global	 Fragmentation, need to build for multiple platforms, with time and costs Managing multiple releases Client-side changes Need to submit app to some stores for approval
n mobile ted and	 Less functionality, unable to use advanced phone features such as camera, global positioning system Small display size Low text input and low bandwidth

SECTION 5

MONITORING AND EVALUATION

Monitoring is the routine tracking of an intervention's performance using data collected on a regular and ongoing basis on specified indicators. This information is used to assess the extent to which an intervention is achieving its intended targets on time and on budget.

Evaluation is an episodic assessment of either a completed or an ongoing programme or intervention to determine the extent to which it achieved its stated objectives efficiently and effectively.

The aims of both monitoring and evaluation are to formulate policies, set priorities, plan, design similar: to provide information to inform decisions, and implement mHypertension projects, change improve outcomes and achieve objectives. the programme in a timely manner, if needed, and allocate the resources to do so.

While monitoring routinely gives information on where a project is at any given time relative to its targets for implementation, an evaluation gives evidence of *why* targets and outcomes are or are not being achieved or the extent to which changes can be attributed to the intervention. The data to be collected for monitoring during a project should be part of the planning process.

Why do we need to monitor and evaluate mHypertension?

There is a need for robust evaluations on the benefits and risks of implementing mHypertension at scale, and its cost-effectiveness. Assessing the processes and contextual factors that influence the implementation of mHypertension programmes will help countries' future efforts, as well as helping others understand how they work, which target group they are best suited to and under which circumstances.

As a guide for planning the evaluation process, a framework or logic model (see Figure 7) that may be expanded and adjusted to the context of the programme and the aspects that are considered key objectives and goals for the programme. For the purposes of simplifying the different interrelated areas of mHealth programmes, this logic model has been divided into two domains -The data from monitoring and evaluation will the person-centred domain and the programmeprovide donors, policy-makers and other decisioncentred domain. An example mHypertension makers at all levels with the relevant information logic model is provided in Figure 8.

5.1 DEVELOPING A MONITORING AND EVALUATION PLAN

Planning for monitoring and evaluation should be part of the mHypertension programme development and implemented from the beginning of the programme. It is especially important to collect baseline data, either before or in the early stages of programme launch, as this will allow for pre-post analysis of programme effectiveness.

FIGURE 7. LOGIC MODEL: FRAMEWORK FOR MONITORING AND EVALUATION OF mHEALTH PROGRAMMES AT SCALE

	INPUT	OUTPUT	OUTCOME	IMPACT
PERSON- CENTERED DOMAIN.	 Outgoing messages Incoming messages Surveys, interviews 	 Reach and registration Information about the user population Ease of understanding messages 	 Improved literacy/ knowledge/outreach Behaviour change Return on investment Technology performance 	Improved health outcomeImproved use of resources
	INPUT	OUTPUT	OUTCOME	IMPACT
PROGRAMME- CENTERED DOMAIN.	 Governance Policy data Resources (finance, HR, ICT architecture) Content development Outreach and promotion Data from "Person- centered domain" 	 Coverage of intervention Intervention quality Interoperability 	 Integration with health systems Improved health literacy Access to intervention 	 Improved health outcomes (SDG 3) Improved digital capacity (SDG 9) Efficiency and efficacy

HR: human resources; ICT: information and communications technology SDG: United Nations Sustainable Development Goal.

FIGURE 8. EXAMPLE LOGIC MODEL FOR mHYPERTENSION EVALUATION

INPUT	ACTIVITIES	OUTPUT	OUTCOME	IMPACT
• Country develops mHTN programme (staff, resources, time, funding, relationship and training with service providers, health centres established)	 Messages are adapted from the mHTN handbook for the local context Messages are pretested with the target audience and finalized Programme recruiters are trained Promotion for the programme conducted SMS fully set up with service provider 	 # of programme recruiters trained # of users enrolled # of messages sent to users # of press releases and amount of media attention 	 Users are satisfied with the programme Messages are received, read, and understood by target population Improvement in knowledge, perceptions, attitudes, intentions and behaviours for each recommended behaviour Health-care providers are satisfied with the programme Improved adherence and compliance to medication Improved awareness of dangers of hypertension and links with heart disease and stroke 	 Users have blood pressure below 140/90 mmHg Fewer users diagnosed with cardiovascular disease
P	ROCESS EVALUATIO	OUTCOME	EVALUATION	

Table 8 lists suggested steps for planning the monitoring and evaluation of an mHealth programme.

TABLE 8. MONITORING AND EVALUATION STEPS AND CONSIDERATIONS

1) Determine the purpose of monitoring and evaluationI Does the mHypertension plate Are the messages easy for us How effective is the marketin How many people from the the How many stayed for the whe Are users improving their known mHealth application? Are user 2) Select indicators*Output I Reach and registration I Information about the user p Ease of understanding the m2) Select indicators*Output I Reach and registration I Information about the user p Ease of understanding the m3) Develop a monitoring and evaluation planWhat is to be monitored and I What is to be monitored and I What are the activities needer fundraising, etc.)? I Who is responsible for monitor I Who will monitoring and evalused? I How will user requests for data fatigue? I Wow will user requests for data fatigue?4) Collect baseline dataDemographic data I Must relate directly to the inci in the monitoring and evalua behaviour, theoretical constribution, theoretical	STEPS	KEY CONSIDERATIONS
 2) Output Select indicators* Reach and registration Information about the user p Ease of understanding the m Outcome Improved literacy/knowledg Behaviour change Return on investment Technology performance 3) What is to be monitored and What is to be monitored and What is to be monitored and What are the activities needer fundraising, etc.)? Who is responsible for monitor When will monitoring and evaluation plan How will user requests for data fatigue? What resources and approval security)? How will data be analysed an Must relate directly to the individual behaviour, theoretical constring readings) 	1) Determine the purpose of monitoring and evaluation	 Does the mHypertension plan Are the messages easy for us How effective is the marketin How many people from the the many stayed for the wh Are users improving their known mHealth application? Are users
 3) Develop a monitoring and evaluation plan What is to be monitored and What are the activities needer fundraising, etc.)? Who is responsible for monitor When will monitoring and evaluated? How will user requests for data fatigue? What resources and approval security)? How will data be analysed an Minimal collection of socioded gender, ethnicity, socioecond Baseline data Must relate directly to the individual behaviour, theoretical constring and evaluated behaviour, theoretical constringed 	2) Select indicators*	Output Reach and registration Information about the user p Ease of understanding the m Outcome Improved literacy/knowledg Behaviour change Return on investment Technology performance
 4) Demographic data Collect baseline data Minimal collection of sociode gender, ethnicity, socioecond Baseline data Must relate directly to the indin the monitoring and evaluate behaviour, theoretical construction readings) 	3) Develop a monitoring and evaluation plan	 What is to be monitored and What are the activities needer fundraising, etc.)? Who is responsible for monitor When will monitoring and evalused? How will user requests for data fatigue? What resources and approval security)? How will data be analysed and
	4) Collect baseline data	 Demographic data Minimal collection of sociode gender, ethnicity, socioecond Baseline data Must relate directly to the ind in the monitoring and evaluat behaviour, theoretical constru- readings)

*Data collection and processing require resources. It is recommended to focus on the set of indicators that are important, feasible and cost-effective to collect, process and analyse.

- tform operate at optimum level?
- sers to understand?
- ng campaign in reaching new users?
- target population did the programme reach? nole duration of the programme?
- owledge, behaviour or attitudes due to the
- ers increasing their use of health services?

opulation essages

je/outreach

l evaluated? ed to monitor and evaluate (training, staffing,

- oring and evaluation activities?
- aluation take place?
- uation be carried out? What measures will be

ata be minimized to prevent user data overload/

Is are required and from whom (e.g. ethics, data

nd distributed?

emographic characteristics, such as age, omic status and other factors

dicators that have been previously identified tion plan (e.g. physical activity and eating ructs such as self-efficacy, blood pressure

5.2 SPECIFIC DATA COLLECTION METHODS

Monitoring

- Recommend setting up a routine reporting mechanism to monitor the core indicators and the key deliverables. Countries with webbased platforms may present the monitoring report as a data dashboard. Determine which report indicators can be automated, the frequency of data collection and the format for analysing the data.
- Decide who will use the monitoring reports (e.g. programme managers, stakeholders) and reflect on the questions they need to have

answered, what decisions they make and how much time they can commit to reviewing the numbers.

Managers should regularly review the progress of participation, technical errors and message activity. In the beginning of a programme, daily monitoring of outgoing messages is recommended. Then weekly or monthly monitoring should be conducted and timed around activities and outputs.

Process evaluation: the periodic assessment of the implementation of a programme in relation to planned activities and overall objectives.

Outcome evaluation: a type of evaluation that is concerned with determining if, and by how much, a programme's activities achieved their intended targets.

Evaluation

While the process evaluation (monitoring) should be done regularly (weekly or monthly), the outcome evaluation can be conducted pre-intervention or post-intervention. For example, baseline data can be collected before launching mHypertension (pre-intervention), and the same measures can be applied at various intervals, such as three months and six months post-launch (post-intervention). To understand long-term effects, another assessment is recommended six months to one year after the mHypertension programme finishes. A simple way to collect data from users is to use the existing mHypertension platform. For instance, survey questions with simple yes/no or numerical answers can be sent to mHypertension users at regular intervals, such as before and after each "module" of text messages. The immediate delivery of survey questions by mobile phones can improve reliability of responses. For further clarification see Figure 9 and refer to Annex 5 for survey question ideas.

Mobile phone survey question 1:

In the past week, on how many days did you do physical activity for 20–30 minutes? Respond with a number from 0 to 7.

FIGURE 9. EXAMPLE QUESTION AND ADMINISTRATION OF A SURVEY QUESTION DELIVERED BY MOBILE PHONE



Table 9 lists different types of evaluation method, along with key considerations for each method.

TABLE 9. METHODS AND DATA FOR mHEALTH EVALUATION

METHOD	PURPOSE	CONSIDERATIONS AND EXAMPLES
Survey	Assess the perceptions, behaviours, knowledge and attitudes and intentions of registered users	 Can be administered on the Web, on a mobile device (recommended), or by an interviewer over the phone or in person Should be short as possible and employ multiple-choice yes/ no questions wherever possible (see Annex 5 for examples)
Focus group	Deepen understanding of users' experiences and explore the factors that may affect the user experience, such as language ability, gender, age, geographical location, etc.	 Conducted by a facilitator in groups of 5-8 people Can also be used to brainstorm ideas for increasing registration, improving the programme and enhancing desired outcomes
Interview	Interviews with users: qualitative feedback on their experience, perceptions and satisfaction with the mHypertension programme Interviews with programme implementers: information on perceived strengths, weaknesses and needs related to programme delivery, processes, staff and management structures, capacity and communications methods	For qualitative methodology of focus groups and interviews, refer to (46)

Objective measure of outcomes	Collect data on biomechanical or physiological outcomes of performing behaviours, often collected in real time		Literature searches can help identify objective assessment methods for relevant health behaviours Examples include devices such as accelerometers or pedometers to measure physical activity, and saliva cotinine test to prove smoking cessation Appointments made and held Clinical markers and biomarkers, such as blood pressure readings, weight and hospitalization for further progression of disease (e.g. cardiovascular disease) If it is not feasible to collect objective measures for all participants, the measures could be used on a subsample of the mHypertension user group, which can be generalized to the total sample after demographic matching
Cost analysis	Compare the cost per user of the mHealth approach with traditional approaches to reaching people in similar hypertension programmes	1	Collect costs on items such as development/planning, staff, marketing and technology

Figure 8 above is an example of a logic model for process and outcome evaluations of mHypertension. Stakeholders in each participating country should be consulted when defining potential evaluation guestions and identify relevant programme outcomes and measures. Other logic models are available (47, 48, 49).

5.3 DISSEMINATION

Once monitoring and evaluation activities and reports are completed, dissemination should occur through the appropriate channels. The list below describes some key considerations for dissemination:

Findings should be disseminated in an audience-friendly and timely manner through:

- formal and informal networks via meetings, newsletters and other forums
- local newspapers and radio programmes
- professional conferences via discussion papers or posters
- □ journals (professional or lay)*
- □ electronic media, such as webpages and email.

* Note that it can be more effective to publish the evaluation results in an academic journal prior to releasing results to the public. Greater confidence can be placed in the results of the mHypertension programme once the findings have been through a rigorous peer-review process.

Conclusion

This handbook provides guidance for the development and implementation of an mHealth programme to support people with hypertension in achieving better control. It describes how an mHypertension programme can be used to supplement existing national hypertension control programmes, and illustrates the steps required for successful implementation. The main components are: operations management, content development and adaptation, promotion and recruitment, technology specification and monitoring and evaluation. The content of the mHypertension programme will complement existing health-care services and routine care offered by health-care professionals. All content in this handbook is based on WHO guidelines, existing research evidence on effectiveness, content and delivery and expert opinions. The templates showcased here should be considered as examples and must be adapted to local context of each participating country.

Resources and Recommended Reading

WHO guideline: recommendations on digital interventions for health system strengthening. Geneva: World Health Organization; 2019 (Licence: CC BY-NC-SA 3.0 IGO; https://apps.who.int/iris/ handle/10665/311980, accessed 1 May 2019).

References

- mHealth: new horizons for health through mobile technologies: second global survey on eHealth. Geneva: World Health Organization; 2011 (http://www.who.int/goe/ publications/goe_mhealth_web.pdf, accessed 1 May 2019).
- A global brief on hypertension. Geneva: World Health Organization; 2013 (http://apps.who.int/iris/ bitstream/10665/79059/1/WHO_DCO_WHD_2013.2_eng.pdf, accessed 1 May 2019).
- Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. Lancet. 2012;380(9859):2224–60. doi:10.1016/S0140-6736(12)61766-8.
- Effect of reduced sodium intake on blood pressure and potential adverse effects in children. Geneva: World Health Organization; 2012 (http://apps.who.int/iris/ bitstream/10665/79328/1/9789241504898_eng.pdf, accessed 1 May 2019).
- HEARTS technical package for cardiovascular disease. In: Cardiovascular disease [website]. Geneva: World Health Organization; 2019 (http://www.who.int/cardiovascular_ diseases/hearts/en/, accessed 1 May 2019).
- Measuring the information society report 2016. Geneva: International Telecommunication Union; 2016 (https:// www.itu.int/en/ITU-D/Statistics/Documents/publications/ misr2016/MISR2016-w4.pdf, accessed 1 May 2019).
- Global status report on noncommunicable diseases, 2010. Geneva: World Health Organization; 2011 (http://www.who. int/nmh/publications/ncd_report_full_en.pdf, accessed 1 May 2019).
- Pfaeffli Dale L, Dobson R, Whittaker R, Maddison R. The effectiveness of mobile-health behaviour change interventions for cardiovascular disease self-management: a systematic review. Eur J Prev Cardiol. 2015; Online first 21 October 2015. doi:10.1177/2047487315613462.
- Chow CK, Islam SM, Farmer A, Bobrow K, Maddison R, Whittaker R et al. Text2PreventCVD: protocol for a systematic review and individual participant data meta-analysis of text message-based interventions for the prevention of cardiovascular diseases. BMJ open. 2016;6(10):e012723.
- Rehman H, Kamal AK, Morris PB, Sayani S, Merchant AT, Virani SS. Mobile health (mHealth) technology for the management of hypertension and hyperlipidemia: slow start but loads of potential. Curr Atheroscler Rep. 2017;19(3):12.
- Anglada-Martinez H, Riu-Viladoms G, Martin-Conde M, Rovira-Illamola M, Sotoca-Momblona JM, Codina-Jane C. Does mHealth increase adherence to medication? Results of a systematic review. Int J Clin Pract. 2015;69(1):9–32.
- Müller AM, Khoo S, Morris T. Text messaging for exercise promotion in older adults from an upper-middle-income country: randomized controlled trial. J Med Internet Res. 2016;18(1).
- King AC, Hekler EB, Grieco LA, Winter SJ, Sheats JL, Buman MP et al. Harnessing different motivational frames via mobile phones to promote daily physical activity and reduce sedentary behavior in aging adults. PloS One. 2013;8(4):e62613

- Eyles H, McLean R, Neal B, Jiang Y, Doughty RN, McLean R et al. A salt-reduction smartphone app supports lower-salt food purchases for people with cardiovascular disease: findings from the SaltSwitch randomised controlled trial. Eur J Prev Cardiol. 2017;(13):1435-44.
- McManus RJ, Mant J, Bray EP, Holder R, Jones MI, Greenfield S et al. (2010). Telemonitoring and self-management in the control of hypertension (TASMINH2): a randomised controlled trial. Lancet. 2010;376(9736):163–72. doi:10.1016/S0140-6736(10)60964-6.
- Park LG, Beatty A, Stafford Z, Whooley MA. Mobile phone interventions for the secondary prevention of cardiovascular disease. Prog Cardiovasc Dis. 2016;58(6):639–50.
- de Silva D. Helping people help themselves: a review of the evidence considering whether it is worthwhile to support self-management. London: Health Foundation; 2011.
- Joe J, Demiris G. Older adults and mobile phones for health: a review. J Biomed Inform. 2013;46(5):947–54.
- Chung SY, Nahm ES. Testing reliability and validity of the eHealth Literacy Scale (eHEALS) for older adults recruited online. Comput Inform Nurs. 2015;33(4):150.
- Rogers EM. Diffusion of innovation. The Free Press: New York; 1983.
- Abroms LC, Whittaker R, Free C, Van Alstyne JM, Schindler-Ruwisch JM. Developing and pretesting a text messaging programme for health behavior change: recommended steps. JMIR mHealth and uHealth. 2015;3(4):e107.
- Hardeman W, Sutton S, Griffin S, Johnston M, White A, Wareham NJ et al. A causal modelling approach to the development of theory-based behaviour change programmes for trial evaluation. Health Educ Res. 2005;20(6):676–87.
- U.S. Department of Health and Human Services. Physical activity and health: a report of the Surgeon General. Atlanta (GA): U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion; 1996 (https://www.cdc.gov/nccdphp/sgr/pdf/chap3.pdf, accessed 1 May 2019).
- Michie S, Richardson M, Johnston M, Abraham C, Francis J, Hardeman W et al. The behavior change technique taxonomy (v1) of 93 hierarchically clustered techniques: building an international consensus for the reporting of behavior change interventions. Ann Behav Med. 2013;46(1):81–95.
- Duff OM, Walsh DM, Furlong BA, O'Connor NE, Moran KA, Woods CB. Behavior change techniques in physical activity ehealth interventions for people with cardiovascular disease: systematic review. J Med Internet Res. 2017 Aug;19(8).
- 26. Webb T, Joseph J, Yardley L, Michie S. Using the internet to promote health behavior change: a systematic review and meta-analysis of the impact of theoretical basis, use of behavior change techniques, and mode of delivery on efficacy. J Med Internet Res. 2010;12(1):e4.
- Cole-Lewis H, Kershaw T. Text messaging as a tool for behavior change in disease prevention and management. Epidemiol Rev. 2010:mxq004.

- Hall AK, Cole-Lewis H, Bernhardt JM. Mobile text messaging for health: a systematic review of reviews. Annu Rev Public Health. 2015;36(1):393–415.
- Buchholz SW, Wilbur J, Ingram D, Fogg L. Physical activity text messaging interventions in adults: a systematic review. Worldviews Evid Based Nurs. 2013;10(3):163-73.
- Free C et al. Smoking cessation support delivered via mobile phone text messaging (txt2stop): a single-blind, randomised trial. Lancet. 2011;378(9785):49-55.
- Maddison R, Pfaeffli L, Whittaker R, Stewart R, Kerr A, Jiang Y, et al. A mobile phone intervention increases physical activity in people with cardiovascular disease: Results from the HEART randomized controlled trial. Eur J Prev Cardiol. 2015;22(6):701-9.
- Pfaeffli Dale L, Whittaker R, Jiang Y, Stewart R, Rolleston A, Maddison R. Text message and internet support for coronary heart disease self-management: results from the Text4Heart randomized controlled trial. J Med Internet Res. 2015;17(10):e237.
- Evans W, Nielsen PE, Szekely DR, Bihm JW, Murray EA, Snider J et al. Dose-response effects of the text4baby mobile health programme: randomized controlled trial. JMIR mHealth and uHealth. 2015;3(1):e12.
- Head KJ, Noar SM, Iannarino NT, Grant Harrington N. Efficacy of text messaging-based interventions for health promotion: a meta-analysis. Soc Sci Med. 2013;97(0):41–8.
- Maar MA, Yeates K, Toth Z, Barron M, Boesch L, Hua-Stewart D et al. Unpacking the black box: a formative research approach to the development of theory-driven, evidencebased, and culturally safe text messages in mobile health interventions. JMIR mHealth and uHealth. 2016;4(1):e10.
- Nigg C, Long C. A systematic review of single health behavior change interventions vs. multiple health behavior change interventions among older adults. Behav Med Pract Policy Res. 2012;2(2):163–79.
- James E, Freund M, Booth A, Duncan MJ, Johnson N, Short CE et al. Comparative efficacy of simultaneous versus sequential multiple health behavior change interventions among adults: a systematic review of randomised trials. Prev Med. 2016;89:211–23.
- Déglise C, Suggs LS, Odermatt P. SMS for disease control: a systematic review of applications in developing countries. J Telemed Telecare. 2012;18(5):273–81.
- Déglise C, Suggs LS, Odermatt P. Short message service (SMS) applications for disease prevention in developing countries. J Med Internet Res. 2012;14(1):e3. doi:10.2196/ jmir.1823.
- Muench F, van Stolk-Cooke K, Morgenstern J, Kuerbis AN, Markle K. Understanding messaging preferences to inform development of mobile goal-directed behavioral interventions. J Med Internet Res. 2014;16(2):e14.
- Notthoff N, Carstensen L. Positive messaging promotes walking in older adults. Psychol Aging. 2014;29(2):329.
- Leroy G, Endicott JE, Kauchak D, Mouradi O, Just M. User evaluation of the effects of a text simplification algorithm using term familiarity on perception, understanding, learning, and information retention. J Med Internet Res 2013;15(7):e144.

- Diet, nutrition and the prevention of chronic diseases: report of a joint WHO/FAO expert consultation, Geneva, 28 January–1 February 2002. Geneva: World Health Organization; 2003 (WHO Technical Report Series, No. 916; https://apps.who.int/iris/bitstream/ handle/10665/42665/WHO_TRS_916.pdf;jsessionid =1488DF1B352833BFDA627AAC245ED390?sequence=1, accessed 1 May 2019).
- Murray CJ, Lauer JA, Hutubessy RC, Niessen L, Tomijima N, Rodgers A et al. Effectiveness and costs of interventions to lower systolic blood pressure and cholesterol: a global and regional analysis on reduction of cardiovascular-disease risk. Lancet. 2003;361(9359):717–25. doi:10.1016/S0140-6736(03)12655-4.
- Powles J, Fahimi S, Micha R, Khatibzadeh S, Shi P, Ezzati M et al. Global, regional and national sodium intakes in 1990 and 2010: a systematic analysis of 24 h urinary sodium excretion and dietary surveys worldwide. BMJ Open. 2013;3(12):e003733-e003733. doi:10.1136/ bmjopen-2013-003733.
- Patton MQ. Enhancing the quality and credibility of qualitative analysis. Health Serv Res. 1999;34(5):1189-1208.
- Ontario Agency for Health Protection and Promotion (Public Health Ontario), Abdi S, Mensah G. Focus on: logic models – a planning and evaluation tool. Toronto (ON): Queen's Printer for Ontario; 2016.
- Logic model guides and examples. In: Toolkits for K4Health [website]. Baltimore (MD): Knowledge for Health Project; 2019 (https://www.k4health.org/toolkits/measuringsuccess/logic-model-guides-and-examples, accessed 1 May 2019).
- Logic models. In: CDC evaluation documents, workbooks and tools [website]. Atlanta (GE): Centers for Disease Control and Prevention; 2019 (https://www.cdc.gov/eval/tools/ logic_models/, accessed 1 May 2019).

ANNEX 1

Examples of mHypertension programmes at the individual and system level

mHypertension Solution Snapshot HIGH BLOOD CARDIOVASCULAR DISEASE **GENERAL POPULATION PRESSURE + RISK** AND OTHER COMPLICATIONS FACTORS Screening and management Prevention Management Message database for health Message database for early Message database for management, screening prompts management, medication promotion SMS adherence, appointment reminders, follow-up reminders Wellness apps for health Risk score apps, medication Monitoring and medication promotion, workplace and stockpiles, workforce training adherence, self-management apps community-based apps APPS Data collection tools HCW electronic decision support tools OTHER Cloud storage, digital health platforms, linking to electronic records mHypertension handbook to DOCUMENTS detail implementation



	ACTIVITIES	SOLUTION TYPE	OBJECTIVE	TOOL	SPECIFIC TARGETS	SOLUTION CONSIDERATIONS	SPECIFIC EXAMPLES	BARRIERS TO CONSIDER	RELATED RESEARCH
H PROMOTION	General health promotion	Individual	Health promotion campaign, encouraging healthy eating habits, salt reduction	SMS, apps, social media	Homemakers	Salt tracking, alternative food choices, recipes and suggestions, guidelines, sharing healthy cooking practices, healthy food discount policy	NHS change4 life campaign, swaps campaign, SaltSwitch, FoodSwitch, WikiHealth, DASH diet	Industry, keeping high engagement, availability and acceptability of alternatives, partnering with health insurers	m-WELLCARE (Shah Ebrahim)
	General health promotion	Individual	Health promotion campaign, encouraging healthy eating habits, salt reduction	SMS, apps, social media	Young people	Share pictures for rewards, games, salt tracker, education and other info		Targeting schools	UDAY study: comprehensive community-based hypertension and diabetes prevention programme
NTION AND HEALI	General health promotion	Individual	Comprehensive health resources to encourage healthy lifestyles	SMS, apps, social media	General population, focus on communities	Availability of health info, online personal health coach	NHS Choices, Arogya World, MeYouHealth leveraging communities, Friendship Bench, mental health support app, Ground Miles, Masiluleke, SMS support messages, Anjna Patient Education		Mama platform
PREVEN	General health promotion	Individual	Comprehensive health resources to encourage healthy lifestyles	SMS, apps, social media	Workplace	Incentive and follow up, bud- dy-matching based on interests and risks, activity levels and monitoring		Partner with health insurers	
	General health promotion	Individual	Informed self-monitoring and tracking		Self-selected	Tracking apps and cloud-based ser- vices, online personal health coach	Fitbit, Runkeeper, Strava, Stepjockey	Wearables integration	
	Surveillance	System	Dashboard with data	Cloud-based, SMS, apps		Dashboard with data, monitoring by local community	Citymapper		
SIS	Monitoring	Individual	Knowing risk score	Apps, devices			Health360		
ND DIAGNO	Screening tools	HCW	Electronic decision support	Apps, devices	нсw	CHW screening toolkit, integrated patient-centred care platform with cloud-based storage	Medic Mobile, SMART health		https://www. ahajournals.org/ doi/full/10.1161/ JAHA.114.001213
ENING /	Training	HCW	Opportunities for networking and experience sharing	Apps, devices	НСЖ	Social sharing platform	Doximity, Sermo, The Rounds		
SCRE	Access to HS	System	Improving access to health system	Apps, devices		Remote consultations	meradoctor.com, Babylon Health, HealthTap		
	Access to HS	System	Improving access to health system	SMS, apps, social media	For individuals	Tool to compare and buy health insurance	HealthSherpa, Kilimo Salama (microin- surance schemes)		
	Management	Individual	Improving management of hyper- tension	SMS, apps, social media		Medication adherence reminders	StAR trial for medication adherence		StAR SMS adher- ence trial
F	Management	Individual	Improving management of hyper- tension	SMS, apps, social media		Self-monitoring tools, medication adherence reminders	Flo (SMS based telehealth), Wellframe, Omada Health, Alere healthcare, Self- Care People, PatientsLikeMe, Propeller Health		
SEMEN'	Management	HCW	Improving diagnosis and man- agement	SMS, apps, social media		Clinical decision-making tools	SnapDx		
MANAQ	Access to medication	System	Improving drug procurement	Cloud-based, SMS, apps	Pharmacists, providers	Central system to aid in tracking and procurement, stock lists of drugs	Central procurement for drugs (<u>http://eaushadhi.rajasthan.gov.in/DWH/start-up/loginAction</u>), e-Aushadhi		
	Management	System	Improving consultation, tracking, history	Cloud-based, SMS, apps	нсw	Paperless records for health care workers	eRecords		
	Management	HCW	Training HCW	Task-shifting for health workers		SnapDx			
	Surveillance	System	Dashboard with data	Cloud-based, SMS, apps		Mapping for hypertension hotspots, dashboard with data, monitoring by local community	Citymapper, Medic Mobile		

ANNEX 2

List of behaviour change techniques included in the mHypertension programme

CONSTRUCT	MICHIE BCT [®]	BCT NO.
Information	Goal setting (behaviour)	1.1
	Information about health consequences	5.1
	Information about emotional consequences	5.6
	Credible source	9.1
Reminders	Prompts/cues	7.1
	Remove aversive stimulus	7.5
Instruction	Instruction on how to perform a behaviour	4.1
	Graded tasks	8.7
Self-management	Goal setting (behaviour)	1.1
	Problem solving	1.2
	Action planning	1.4
	Self-monitoring of behaviour	2.3
	Social support (unspecified)	3.1
	Social support (practical)	3.2
	Social support (emotional)	3.3
	Monitoring of emotional consequences	5.4
	Behaviour substitution	8.2
	Verbal persuasion about capability	15.1

^a Michie S, Richardson M, Johnston M, Abraham C, Francis J, Hardeman W et al. The behavior change technique taxonomy (v1) of 93 hierarchically clustered techniques: building an international consensus for the reporting of behavior change interventions. Ann Behav Med. 2013;46(1):81-95. BCT: behaviour change technique.

ANNEX 3

The mHypertension Message Library is available upon request. To request the mHypertension message library, please contact:

BHBM@who.int or Pujaris@who.int

ANNEX 4

Examples of mHypertension programme algorithms

During the content development phase, it is important to test different programme algorithms with the end users. An important consideration is whether modules should be delivered simultaneously, as this provides variety and reminds users about the different lifestyle changes needed throughout the programme, or sequentially, as some users may find this overwhelming and may want to concentrate on one behaviour at a time. Here are two example algorithms, highlighting how the different mHypertension modules can be presented simultaneously or sequentially.

EXAMPLE 1. mHYPERTENSION SIMULTANEOUS BEHAVIOUR ALGORITHM

An mHypertension user has selected all five modules in simultaneous format. User receives one message from each module per week.

WEEK	DAY	MODULE	EXAMPLE MESSAGE
1	Mon	Adherence	Hello [name], welcome to mHypertension! You will start to receive supportive text messages about healthy living to control your blood pressure
	Tue	Healthy eating	You can control your blood pressure by using less salt in your food. Protect your heart
	Wed	Physical activity	Regular physical activity, like a brisk walk each day, will help control your blood pressure and protect you against heart attack and stroke
	Thu		
	Fri	Tobacco cessation	Stopping tobacco use can reduce your risk of heart attack, stroke, cancer and other diseases. There is no "safe" amount of tobacco
	Sat	Alcohol	Limiting alcohol intake can reduce your risk of heart attack, stroke, cancer and other diseases. Aim for less than 2 drinks per day
	Sun		
2-23	Mon	Adherence	If your blood pressure is high, you have likely been prescribed medication. Taking it will help control your blood pressure and keep your heart healthy
	Tue	Healthy eating	To stay healthy, you ideally need to eat at least 5 servings of fruit or vegetables a day. Try a rainbow of colours each day
	Wed	Physical activity	Regular exercise like walking, gardening, dancing, or playing a sport helps keep your mind and body strong. Make it a daily habit for good health!
	Thu		
	Fri	Tobacco cessation	Quitting tobacco is the most important thing you can do for your health and to reduce your risk of heart disease
	Sat	Alcohol	Avoid alcohol to control your blood pressure and reduce risk of heart disease. Remember that less is best!

VEEK	DAY	MODULE	EXAMPLE MESS
	Sun		
4	Mon	Adherence	Keep on top of pressure should doctor
	Tue	Healthy eating	Stock cubes and to avoid them
	Wed	Physical activity	Change your ha that you normal physical activity
	Thu		
	Fri	Tobacco cessation	Think about wh smoking. Stay f
	Sat	Alcohol	Try a quick walk be rewarded tw feel good too
	Sun	Closing	Well done! You' focused, you sh

2

EXAMPLE 2. mHYPERTENSION SEQUENTIAL BEHAVIOUR ALGORITHM

A non-smoking mHypertension user has selected four modules in sequential format. The user will receive four messages per week from one module at a time. Note that, if a user selects the tobacco cessation and/or alcohol modules, the number of messages per week, the number of weeks per module, and the total length of the mHypertension programme may vary.

WEEK	DAY	MODULE	EXAMPLE MES
1-6	Mon	Adherence	Hello [name], v receive suppor control your bl
	Tue		
	Wed	Adherence	If your blood p medication. Ta and keep your
	Thu		
	Fri	Adherence	Controlling you stroke. Check y needed lifestyl
	Sat	Adherence	Your blood pre taking it even i
	Sun		
7-12	Mon	Healthy eating	You can help c your food. Prot
	Tue		
	Wed	Healthy eating	To stay healthy fruit or vegetab
	Thu		
	Fri	Healthy eating	Prepare food u family maintain off the table!

AGE

- regular blood pressure screening. Your blood be less than 140/90. If it high, see your
- d ready-made soups contain a lot of salt. Try
- bits for healthy living! Try swapping time ly spend sitting down, like watching TV, with
- at you are gaining and why you want to quit ocused. It will get easier
- if you feel the urge to drink. Your health will vo times over! Physical activity can make you
- ve finished the programme. If you stay ould gain further benefits

SAGE

- velcome to mHypertension! You will start to tive text messages about healthy living to ood pressure
- ressure is high, you have likely been prescribed king it will help control your blood pressure heart healthy
- or blood pressure can prevent heart attack or our blood pressure regularly and make any e changes
- ssure medication is for the long term. Keep f you feel well
- ontrol your blood pressure by using less salt in ect your heart
- you ideally need to eat at least 5 servings of ples a day. Try a rainbow of colours each day
- sing only a little bit of salt to help you and your normal blood pressure. Keep the salt shaker

WEEK	DAY	MODULE	EXAMPLE MESSAGE
	Sat	Healthy eating	An adult only needs half a teaspoon of salt per day (less than 5 g, or 2 g of sodium). Look for salt and/or sodium on nutrition facts food labels
	Sun		
13-18	Mon	Physical activity	Regular physical activity, like a brisk walk each day, will help control your blood pressure and protect you against heart attack or stroke
	Tue		
	Wed	Physical activity	Regular exercise like walking, gardening, dancing, or playing a sport helps keep your mind and body strong. Make it a daily habit for good health!
	Thu		
	Fri	Physical activity	Just starting out? See how many minutes you can walk without stopping. Then increase by 1-2 minutes each time
	Sat	Physical activity	Moderate physical activity, like a brisk walk, should make you breathe slightly faster and make your heart beat slightly faster
	Sun		
19-24	Mon	Alcohol	Limiting alcohol intake can reduce your risk of heart attack, stroke, cancer and other diseases. Aim for less than 2 drinks per day
	Tue		
	Wed	Alcohol	Avoid alcohol to control your blood pressure and reduce risk of heart disease. Remember that less is best!
	Thu		
	Fri	Alcohol	The healthiest bet is to avoid alcohol. Aim for less than 2 drinks per day with no alcohol on at least 2 days per week
	Sat	Alcohol	Despite what you may hear in the media, alcohol should not be taken for "health" reasons. Limit your use to less than 2 drinks per day
	Sun	Closing	Well done! You've finished the programme. If you stay focused, you should gain further benefits

ANNEX 5. EXAMPLES OF EVALUATION QUESTIONS

Asking simple questions by text message is a quick and easy way to collect data on outcomes at various stages throughout the programme. To simplify the two-way messaging process, responses should be limited to Yes/No or numeric values. Be specific in the question about the type of response is required. Consider ways to review incoming responses and collate the data.

MODULE	EXAMPLE MESSAGES
Adherence	Have you measured your blood pressure in the last month? Reply with 1 for yes or 2 for no
	Was your last blood pressure reading less than 140/90 mmHg? Reply with 1 for yes or 2 for no
	Did you take your medications yesterday? Reply with 1 for yes or 2 for no
	Did you take your medications each day last week? Reply with 1 for yes or 2 for no
Healthy eating	Have you eaten 5 servings of fruit and vegetables today? Reply with 1 for yes or 2 for no
	Have you started to use less salt in cooking and eating? Reply with 1 for yes or 2 for no
	Have you reduced the amount of fat you cook with and eat? Reply with 1 for yes or 2 for no
Physical activity	Did you do 30 minutes of physical activity today? Reply with 1 for yes or 2 for no
	In the past week, on how many days did you do physical activity for 20-30 minutes? Reply with a number from 0 to 7
Tobacco cessation	Have you quit smoking or using tobacco? Reply with 1 for yes or 2 for no
	Have you been smoke- or tobacco-free for the past 7 days? Reply with 1 for yes or 2 for no
	Have you been smoke or tobacco free for the past 30 days? Reply with 1 for yes or 2 for no
Harmful use of alcohol	Have you had at least 2 alcohol-free days in the last week? Reply with 1 for yes or 2 for no
	Have you had 2 or fewer drinks per day in the last week? Reply with 1 for yes or 2 for no

ISBN 978-92-4-000144-2 (WHO) ISBN 978-92-61-29851-7 (ITU)



Printed in Switzerland Geneva, 2017 Photo credit: Istockphoto/gawrav

International Telecommunication Union

Place des Nations CH-1211 Geneva 20 Switzerland

website: mhealth4ncd.itu.int
website: http://www.who.int/ncds/prevention/be-healthy-be-mobile/en/
e-mail: mhealth@itu.int